

**MATHEMATICAL ENGINEERING
TECHNICAL REPORTS**

**Asymptotic distribution of Wishart matrix for
block-wise dispersion of population eigenvalues**

Yo SHEENA and Akimichi TAKEMURA

METR 2006-49

September 2006

DEPARTMENT OF MATHEMATICAL INFORMATICS
GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY
THE UNIVERSITY OF TOKYO
BUNKYO-KU, TOKYO 113-8656, JAPAN

WWW page: <http://www.i.u-tokyo.ac.jp/mi/mi-e.htm>

The METR technical reports are published as a means to ensure timely dissemination of scholarly and technical work on a non-commercial basis. Copyright and all rights therein are maintained by the authors or by other copyright holders, notwithstanding that they have offered their works here electronically. It is understood that all persons copying this information will adhere to the terms and constraints invoked by each author's copyright. These works may not be reposted without the explicit permission of the copyright holder.

Asymptotic Distribution of Wishart Matrix for Block-wise Dispersion of Population Eigenvalues

Yo Sheena* and Akimichi Takemura†

September, 2006

Abstract

This paper deals with the asymptotic distribution of Wishart matrix and its application to the estimation of the population matrix parameter when the population eigenvalues are block-wise infinitely dispersed. We show that the appropriately normalized eigenvectors and eigenvalues asymptotically generate two Wishart matrices and one normally distributed random matrix, which are mutually independent. For a family of orthogonally equivariant estimators, we calculate the asymptotic risks with respect to the entropy or the quadratic loss function and derive the asymptotically best estimator among the family. We numerically show 1) the convergence in both the distributions and the risks are quick enough for a practical use, 2) the asymptotically best estimator is robust against the deviation of the population eigenvalues from the block-wise infinite dispersion.

Key words and phrases: covariance matrix, Wishart distribution, quadratic loss, Stein's loss, asymptotic risk

1 Introduction

Suppose that a p -dimensional random vector \mathbf{y} has the covariance matrix Σ . The inference for Σ has been studied in enormous amount of literature and is still an important topic from both theoretical and practical points of view. Often we assume some structure of Σ , i.e., restriction on its parameter space $\{\Sigma \mid \Sigma > 0\}$. A structure, in some cases, arises from a theoretical reason behind the data. In other cases, it appears as a result of exploratory analysis such as principle component analysis or exploratory factor analysis.

For example suppose that \mathbf{y} is generated in the following multivariate linear model;

$$\mathbf{y} = \mathbf{B}\mathbf{x} + \mathbf{e}, \quad (1)$$

where \mathbf{B} is a $p \times m$ coefficient (factor loading) matrix with rank $\mathbf{B} = m$, \mathbf{x} is a latent $m \times 1$ random vector (common factor) and $p \times 1$ vector \mathbf{e} is an error term (unique factor) which is independently

*Department of Economics, Shinshu University

†Graduate School of Information Science and Technology, University of Tokyo

distributed from \mathbf{x} . If we further assume that \mathbf{e} has $\sigma^2 \mathbf{I}_p$ (\mathbf{I}_p : p -dimensional identity matrix) as its covariance matrix, $\mathbf{\Sigma}$ is written as

$$\mathbf{\Sigma} = \mathbf{B}\mathbf{\Sigma}_x\mathbf{B}' + \sigma^2\mathbf{I}_p,$$

where $\mathbf{\Sigma}_x$ is the nonsingular covariance matrix of \mathbf{x} . In this case $\mathbf{\Sigma}$ has the eigenvalues $\lambda_1 \geq \dots \geq \lambda_p$ given by

$$\lambda_i = \begin{cases} \tau_i + \sigma^2, & \text{if } i = 1, \dots, m, \\ \sigma^2, & \text{if } i = m + 1, \dots, p. \end{cases} \quad (2)$$

where $\tau_i > 0$, $i = 1, \dots, m$, are the eigenvalues of $\mathbf{B}\mathbf{\Sigma}_x\mathbf{B}'$. It is often observed that σ^2 is quite small compared to τ_i 's, which means that the first group of eigenvalues $(\lambda_1, \dots, \lambda_m)$ is very large compared to the second group $(\lambda_{m+1}, \dots, \lambda_p)$. In this paper we call this state as “(two-)block-wise dispersion” of the population eigenvalues.

What would happen to the sample covariance matrix, when the eigenvalues of population covariance matrix are “infinitely” dispersed? This is an interesting question from a theoretical standpoint. Takemura and Sheena (2005) and Sheena and Takemura (2006) deal with this problem under “total dispersion” of population eigenvalues, namely

$$(\lambda_2/\lambda_1, \lambda_3/\lambda_2, \dots, \lambda_p/\lambda_{p-1}) \rightarrow \mathbf{0}.$$

This paper is a generalization of Takemura and Sheena (2005) from a theoretical point of view, while the practical motivation is as follows; as we saw above, we often come across a practical situation where the population eigenvalues are block-wise dispersed. It is helpful for the inference on $\mathbf{\Sigma}$ in practical situations to understand the behavior of the sample covariance matrix, when the population eigenvalues are block-wise “infinitely” dispersed. The state of the population eigenvalues being infinitely dispersed is a theoretical approximation, but understanding the limiting behavior leads to a better insight on its neighborhood where the eigenvalues are “largely” dispersed.

Now we formally state the framework of this paper. Let $\mathbf{S} = (s_{ij})$ be distributed according to Wishart distribution $\mathbf{W}_p(n, \mathbf{\Sigma})$, where p is the dimension, n is the degrees of freedom, and $\mathbf{\Sigma}$ is the covariance matrix. The spectral decompositions of $\mathbf{\Sigma}$ and \mathbf{S} are given by

$$\mathbf{\Sigma} = \mathbf{\Gamma}\mathbf{\Lambda}\mathbf{\Gamma}', \quad \mathbf{S} = \mathbf{G}\mathbf{L}\mathbf{G}',$$

where $\mathbf{G}, \mathbf{\Gamma} \in \mathcal{O}(p)$, the group of $p \times p$ orthogonal matrices, and $\mathbf{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_p)$, $\mathbf{L} = \text{diag}(l_1, \dots, l_p)$, are diagonal matrices with the eigenvalues $\lambda_1 \geq \dots \geq \lambda_p > 0$, $l_1 \geq \dots \geq l_p > 0$ of $\mathbf{\Sigma}$ and \mathbf{S} , respectively. We use the notations $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_p)$ and $\mathbf{l} = (l_1, \dots, l_p)$ hereafter. By the requirement that

$$\tilde{\mathbf{G}} = (\tilde{g}_{ij}) = \mathbf{\Gamma}'\mathbf{G}$$

has positive diagonal elements, the spectral decomposition $\mathbf{S} = \mathbf{G}\mathbf{L}\mathbf{G}'$ is almost surely uniquely determined. Then almost surely there exists a one-to-one correspondence between the set $\{\mathbf{S} \mid \mathbf{S} > 0\}$ and $\mathcal{L} \times \mathcal{O}^+(p)$, where

$$\mathcal{L} = \{\mathbf{l} \mid l_1 > \dots > l_p > 0\}, \quad \mathcal{O}^+(p) = \{\tilde{\mathbf{G}} \in \mathcal{O}(p) \mid \tilde{g}_{ii} > 0, 1 \leq i \leq p\}.$$

Let m (m_i in Subsection 2.3) denote the dividing point of the first block and the second block of the eigenvalues. Now we parameterize $\boldsymbol{\lambda}, \mathbf{l}$ as follows;

$$\lambda_i = \begin{cases} \xi_i \alpha, & \text{if } i = 1, \dots, m, \\ \xi_i \beta, & \text{if } i = m + 1, \dots, p, \end{cases} \quad (3)$$

$$l_i = \begin{cases} d_i \alpha, & \text{if } i = 1, \dots, m, \\ d_i \beta, & \text{if } i = m + 1, \dots, p, \end{cases} \quad (4)$$

In this paper we always consider ξ 's are given and fixed. We also use the notations,

$$\boldsymbol{\Xi} = \text{diag}(\xi_1, \dots, \xi_p), \quad \boldsymbol{\xi} = (\xi_1, \dots, \xi_p),$$

$$\mathbf{D} = \text{diag}(d_1, \dots, d_p), \quad \mathbf{d} = (d_1, \dots, d_p).$$

We will investigate the asymptotic distribution of \mathbf{S} as β/α goes to 0 while $\boldsymbol{\Xi}$ is fixed and its application to the estimation of $\boldsymbol{\Sigma}$. The state $\beta/\alpha \approx 0$ means that the eigenvalues of $\boldsymbol{\Sigma}$ are two-block-wise ‘‘largely’’ dispersed. In the following, the notation $\beta/\alpha \rightarrow 0$ means a limiting operation $n \rightarrow \infty$ with arbitrary sequences α_n, β_n , $n = 1, 2, \dots$, such that $\beta_n/\alpha_n \rightarrow 0$.

We briefly describe the content of the following sections. In Subsection 2.1 we prepare a local coordinate system of $\mathcal{O}^+(p)$ around \mathbf{I}_p . In Subsection 2.2 we present our main results on asymptotic distributions and we further discuss the case of multi-block-wise infinite dispersion in Subsection 2.3. Section 3 deals with the estimation of $\boldsymbol{\Sigma}$ from decision-theoretic framework. In Subsection 3.1 we introduce orthogonally equivariant estimators and two loss functions and in Subsection 3.2 we calculate the asymptotic risks. We concentrate on the special case of block-wise identity covariance matrices in Subsection 3.3, which is practically important, and we propose the best estimator for the case with respect to each loss function. In Subsection 3.4 the convergence speed of both distributions and risks are numerically evaluated. Together with the application to discriminant analysis, the numerical comparisons show the superiority of the new estimators. In Appendix we present the proofs of two lemmas and discuss analytical calculation of the asymptotic risks.

Before concluding this subsection, we introduce some notational conventions in this paper. In the sections other than Subsection 2.3, we always consider a same two-block partition of matrices. For $\mathbf{A} = (a_{ij})$, a $p \times p$ matrix, \mathbf{A}_{ij} ($1 \leq i, j \leq 2$) denotes the (i, j) -block in the partition

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix}, \quad \mathbf{A}_{11} : m \times m, \quad \mathbf{A}_{22} : (p - m) \times (p - m).$$

If \mathbf{A} is block diagonal, i.e. $\mathbf{A}_{12} = \mathbf{A}_{21} = \mathbf{0}$, we write

$$\mathbf{A} = \text{diag}(\mathbf{A}_{11}, \mathbf{A}_{22}) = \begin{pmatrix} \mathbf{A}_{11} & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_{22} \end{pmatrix}.$$

For the particular case of diagonal matrix $\mathbf{A} = \text{diag}(a_1, \dots, a_p)$, we simply write $\mathbf{A}_1, \mathbf{A}_2$ instead of $\mathbf{A}_{11}, \mathbf{A}_{22}$, i.e. $\mathbf{A}_1 = \text{diag}(a_1, \dots, a_m)$, $\mathbf{A}_2 = \text{diag}(a_{m+1}, \dots, a_p)$. Let $\mathbf{a} = (a_{ij})_{1 \leq j < i \leq p}$ denote the vector of the elements in the lower triangular part of \mathbf{A} , which is correspondingly partitioned as $\mathbf{a} = (\mathbf{a}_{11}, \mathbf{a}_{22}, \mathbf{a}_{21})$, where

$$\mathbf{a}_{11} = (a_{ij})_{1 \leq j < i \leq m}, \quad \mathbf{a}_{22} = (a_{ij})_{m+1 \leq j < i \leq p}, \quad \mathbf{a}_{21} = (a_{ij})_{1 \leq j \leq m < i \leq p}.$$

If \mathbf{a} is a p -dimensional row vector, i.e., $\mathbf{a} = (a_1, \dots, a_p)$, then we make a partition of \mathbf{a} as

$$\mathbf{a} = (\mathbf{a}_1, \mathbf{a}_2), \quad \mathbf{a}_1 = (a_1, \dots, a_m), \quad \mathbf{a}_2 = (a_{m+1}, \dots, a_p).$$

We write $\text{etr } X = \exp(\text{tr } X)$ for a square matrix X .

2 Asymptotic Distribution

2.1 Local Coordinates

We consider a local coordinate of $\mathcal{O}^+(p)$, $\mathbf{u} = (u_{ij})_{1 \leq j < i \leq p}$, around the identity matrix \mathbf{I}_p . For the proof of the existence of such coordinate, see Appendix B of Takemura and Sheena (2005). We have the following open sets C_ϵ, U, V and functions ϕ_{ij} , $1 \leq i \leq j \leq p$;

$$\begin{aligned} C_\epsilon &= \{\mathbf{u} \mid |u_{ij}| < \epsilon, 1 \leq j < i \leq p\} \subset R^{p(p-1)/2}, \\ \mathbf{0} &\in U \subset \bar{U} \subset C_\epsilon, \\ \mathbf{I}_p &\in V \subset \mathcal{O}^+(p), \end{aligned}$$

and $\phi_{ij}(\mathbf{u})$ is a C^∞ function on C_ϵ such that $\mathbf{G}(\mathbf{u}) = (g_{ij}(\mathbf{u}))$ defined by

$$\begin{cases} g_{ij}(\mathbf{u}) = \phi_{ij}(\mathbf{u}), & 1 \leq i \leq j \leq p, \\ g_{ij}(\mathbf{u}) = u_{ij}, & 1 \leq j < i \leq p, \end{cases} \quad (5)$$

is a one-to-one function from U onto V . Using V we can construct a finite open covering of $\mathcal{O}^+(p)$ as follows. For $\mathbf{H}_1 \in \mathcal{O}^+(m)$, $\mathbf{H}_2 \in \mathcal{O}^+(p-m)$, let

$$V(\mathbf{H}_1, \mathbf{H}_2) = \text{diag}(\mathbf{H}_1, \mathbf{H}_2)V \cap \mathcal{O}^+(p) = \{\mathbf{G} \mid \mathbf{G} = \text{diag}(\mathbf{H}_1, \mathbf{H}_2)\mathbf{G}^*, \exists \mathbf{G}^* \in V\} \cap \mathcal{O}^+(p).$$

denote the open neighborhood of $\text{diag}(\mathbf{H}_1, \mathbf{H}_2)$. Let

$$\mathcal{O}(m, p-m) = \{\text{diag}(\mathbf{H}_1, \mathbf{H}_2) \mid \mathbf{H}_1 \in \mathcal{O}^+(m), \mathbf{H}_2 \in \mathcal{O}^+(p-m)\}$$

then

$$\mathcal{O}(m, p-m) \subset \bigcup_{\mathbf{H}_1 \in \mathcal{O}^+(m), \mathbf{H}_2 \in \mathcal{O}^+(p-m)} V(\mathbf{H}_1, \mathbf{H}_2).$$

Since $\mathcal{O}(m, p-m)$ is compact, we can choose a finite number of sets $O^{(\tau)} = V(\mathbf{H}_1^{(\tau)}, \mathbf{H}_2^{(\tau)})$, $\tau = 1, \dots, T$, such that $\bigcup_{\tau=1}^T O^{(\tau)} \supset \mathcal{O}(m, p-m)$. Let $O^{(0)} = \mathcal{O}^+(p) \setminus \mathcal{O}(m, p-m)$, then we have a finite open covering $\{O^{(\tau)}\}_{\tau=0}^T$ of $\mathcal{O}^+(p)$. We denote the partition of unity subordinate to $\{O^{(\tau)}\}_{\tau=0}^T$ by $\{\iota_\tau\}_{\tau=0}^T$. Namely for each τ , ι_τ is a continuous function from $\mathcal{O}^+(p)$ to $[0, 1]$, the support of ι_τ is contained in $O^{(\tau)}$, and $\sum_{\tau=0}^T \iota_\tau(\mathbf{G}) \equiv 1$.

For $O^{(\tau)}$, $1 \leq \tau \leq T$, we can use \mathbf{u} as a local coordinate since \mathbf{G} in $O^{(\tau)}$ can be uniquely expressed as $\mathbf{H}^{(\tau)}\mathbf{G}(\mathbf{u})$ with some \mathbf{u} in U , where

$$\mathbf{H}^{(\tau)} = \text{diag}(\mathbf{H}_1^{(\tau)}, \mathbf{H}_2^{(\tau)}), \quad \tau = 1, \dots, T. \quad (6)$$

As we will see later, we do not need a local coordinate for $O^{(0)}$, since the measure of this area asymptotically vanishes.

Now we have (\mathbf{l}, \mathbf{u}) as a local coordinate on each $\mathcal{L} \times O^{(\tau)}$, $\tau = 1, \dots, T$. We need another local coordinate to investigate the asymptotic behavior of \mathbf{S} . Let $\mathbf{q} = (q_{ij})_{1 \leq j < i \leq p}$ be defined as follows as another coordinate on $O^{(\tau)}$ for a fixed τ , $\tau = 1, \dots, T$; if $1 \leq j \leq m < i \leq p$,

$$\begin{aligned} q_{ij} &= l_j^{1/2} \lambda_i^{-1/2} \sum_{t=m+1}^p (\mathbf{H}_2^{(\tau)})_{i-m, t-m} u_{tj} \\ &= \alpha^{1/2} \beta^{-1/2} d_j^{1/2} \xi_i^{-1/2} \sum_{t=m+1}^p (\mathbf{H}_2^{(\tau)})_{i-m, t-m} u_{tj} \end{aligned} \quad (7)$$

and $q_{ij} = u_{ij}$ otherwise. If we use matrices $\mathbf{Q} = (q_{ij})$, $\mathbf{U} = (u_{ij})$ and their partitions, (7) is the same as

$$\mathbf{Q}_{21} = \alpha^{1/2} \beta^{-1/2} \mathbf{\Xi}_2^{-1/2} \mathbf{H}_2^{(\tau)} \mathbf{U}_{21} \mathbf{D}_1^{1/2}, \quad \mathbf{Q}_{11} = \mathbf{U}_{11}, \quad \mathbf{Q}_{22} = \mathbf{U}_{22}. \quad (8)$$

Conversely

$$\mathbf{U}_{21} = \alpha^{-1/2} \beta^{1/2} \mathbf{H}_2^{(\tau)'} \mathbf{\Xi}_2^{1/2} \mathbf{Q}_{21} \mathbf{D}_1^{-1/2}, \quad \mathbf{U}_{11} = \mathbf{Q}_{11}, \quad \mathbf{U}_{22} = \mathbf{Q}_{22}, \quad (9)$$

or

$$u_{ij} = \begin{cases} \alpha^{-1/2} \beta^{1/2} \sum_{t=m+1}^p (\mathbf{H}_2^{(\tau)})_{t-m, i-m} q_{tj} \xi_t^{1/2} d_j^{-1/2}, & \text{if } 1 \leq j \leq m < i \leq p, \\ q_{ij}, & \text{otherwise.} \end{cases} \quad (10)$$

Pairing $\mathbf{q} = (q_{ij})_{1 \leq j < i \leq p}$ with $\mathbf{d} = (d_1, \dots, d_p)$, we have another local coordinate (\mathbf{d}, \mathbf{q}) on $\mathcal{D} \times O^{(\tau)}$, where

$$\mathcal{D} = (\mathcal{D}_1 \times \mathcal{D}_2) \cap \mathcal{D}_3 \quad (11)$$

with

$$\begin{aligned} \mathcal{D}_1 &= \{\mathbf{d}_1 \mid d_1 > \dots > d_m > 0\} \\ \mathcal{D}_2 &= \{\mathbf{d}_2 \mid d_{m+1} > \dots > d_p > 0\} \\ \mathcal{D}_3 &= \{(\mathbf{d}_1, \mathbf{d}_2) \mid d_m/d_{m+1} > \beta/\alpha\}. \end{aligned}$$

The Jacobian of the transformation $J((\mathbf{l}, \mathbf{u}) \rightarrow (\mathbf{d}, \mathbf{q}))$ is given by

$$\begin{aligned} \left| \det \left(\frac{\partial(\mathbf{l}, \mathbf{u})}{\partial(\mathbf{d}, \mathbf{q})} \right) \right| &= \left| \det \left(\frac{\partial \mathbf{l}}{\partial \mathbf{d}} \right) \right| \left| \det \left(\frac{\partial \mathbf{u}}{\partial \mathbf{q}} \right) \right| \\ &= \alpha^m \beta^{p-m} \prod_{j \leq m < i} \left(d_j^{-\frac{1}{2}} \xi_i^{\frac{1}{2}} \alpha^{-\frac{1}{2}} \beta^{\frac{1}{2}} \right) \\ &= \alpha^{m - \frac{m(p-m)}{2}} \beta^{p-m + \frac{m(p-m)}{2}} \prod_{j=1}^m d_j^{-\frac{(p-m)}{2}} \prod_{i=m+1}^p \xi_i^{\frac{m}{2}}. \end{aligned} \quad (12)$$

2.2 Main Results

The following theorem says that $\widetilde{\mathbf{G}}$ asymptotically separates into two orthogonal matrices $\widetilde{\mathbf{G}}_{11}$, $\widetilde{\mathbf{G}}_{22}$ on the diagonal blocks.

Theorem 1

- 1 As $\beta/\alpha \rightarrow 0$, $\widetilde{\mathbf{G}}_{21} \xrightarrow{p} \mathbf{0}$.
- 2 $\lim_{\beta/\alpha \rightarrow 0} P(\widetilde{\mathbf{G}} \in O) = 1$ for any open set $O \subset \mathcal{O}^+(p)$ including $\mathcal{O}(m, p-m)$.

Proof. Since 2 is easily proved from 1, we only prove 1 here. Let

$$\bar{\mathbf{S}} = (\bar{s}_{ij}) = \mathbf{\Lambda}^{-\frac{1}{2}} \mathbf{\Gamma}' \mathbf{S} \mathbf{\Gamma} \mathbf{\Lambda}^{-\frac{1}{2}} = \mathbf{\Lambda}^{-\frac{1}{2}} \widetilde{\mathbf{G}} \mathbf{L} \widetilde{\mathbf{G}}' \mathbf{\Lambda}^{-\frac{1}{2}} \sim \mathbf{W}_p(n, \mathbf{I}_p),$$

Suppose $1 \leq j \leq m < i \leq p$. Note that

$$\bar{s}_{ii} = (\tilde{g}_{i1}^2 l_1 + \cdots + \tilde{g}_{ip}^2 l_p) \lambda_i^{-1}.$$

Therefore

$$\tilde{g}_{ij}^2 \leq \bar{s}_{ii} \frac{\lambda_i}{l_j} = \bar{s}_{ii} \frac{\lambda_j}{l_j} \frac{\lambda_i}{\lambda_j} \leq \bar{s}_{ii} \frac{\lambda_j}{l_j} \frac{\xi_i}{\xi_j} \frac{\beta}{\alpha}. \quad (13)$$

Since \bar{s}_{ii} is distributed independently of $\mathbf{\Sigma}$, for any $\epsilon > 0$, there exists M such that

$$P(\bar{s}_{ii} < M) > 1 - \epsilon, \quad \forall \mathbf{\Sigma}. \quad (14)$$

Besides, from the result of Lemma 1 of Takemura & Sheena (2005), for any $\epsilon > 0$, there exists C such that

$$P\left(\frac{\lambda_j}{l_j} < C\right) > 1 - \epsilon, \quad \forall \mathbf{\Sigma}. \quad (15)$$

From (14) and (15) we have

$$\bar{s}_{ii} \frac{\lambda_j}{l_j} \frac{\beta}{\alpha} \xrightarrow{p} 0 \quad \text{as} \quad \frac{\beta}{\alpha} \rightarrow 0.$$

From this fact and (13) we have

$$\tilde{g}_{ij}^2 \xrightarrow{p} 0 \quad \text{as} \quad \frac{\beta}{\alpha} \rightarrow 0, \quad 1 \leq \forall j \leq m < \forall i \leq p.$$

■

Next we state a rather technical lemma, which will be used in the proofs of some theorems. Consider a random variable $x(\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta)$. We are often interested in the asymptotic expectation of $x(\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta)$ as $\beta/\alpha \rightarrow 0$ while $\mathbf{\Gamma}$ is fixed. For fixed $\mathbf{\Gamma}$ and $\mathbf{H}^{(\tau)} = \text{diag}(\mathbf{H}_1^{(\tau)}, \mathbf{H}_2^{(\tau)})$, $\mathbf{H}_1^{(\tau)} \in \mathcal{O}^+(m)$, $\mathbf{H}_2^{(\tau)} \in \mathcal{O}^+(p-m)$, somewhat abusing the notation, let

$$x(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \mathbf{\Gamma}, \mathbf{H}^{(\tau)}) = x(\mathbf{\Gamma} \mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)), \mathbf{l}(\mathbf{d}, \alpha, \beta), \boldsymbol{\lambda}(\boldsymbol{\xi}, \alpha, \beta), \alpha, \beta) \quad (16)$$

for emphasizing the right-hand side as the function of $(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)$, where $\mathbf{G}(\mathbf{u})$, $\mathbf{u}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)$, $\mathbf{l}(\mathbf{d}, \alpha, \beta)$, $\boldsymbol{\lambda}(\boldsymbol{\xi}, \alpha, \beta)$ are respectively defined by (5), (10), (4) and (3). For $\mathbf{u} = (\mathbf{u}_{11}, \mathbf{u}_{22}, \mathbf{u}_{21})$, we have

$$\lim_{\beta/\alpha \rightarrow 0} \mathbf{u}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta) = \lim_{\beta/\alpha \rightarrow 0} (\mathbf{u}_{11}(\mathbf{q}_{11}), \mathbf{u}_{22}(\mathbf{q}_{22}), \mathbf{u}_{21}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)) = (\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}), \quad (17)$$

hence

$$\lim_{\beta/\alpha \rightarrow 0} \mathbf{G}(\mathbf{u}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)) = \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}). \quad (18)$$

Lemma 1 Suppose that there exist some $a < 1/2$ and $b > 0$ such that

$$|x(\mathbf{\Gamma}\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta)| \leq b \operatorname{etr}(a\mathbf{G}\mathbf{L}\mathbf{G}'\boldsymbol{\Lambda}^{-1}) \quad \text{a.e. in } (\mathbf{G}, \mathbf{l}) \quad (19)$$

and suppose that for each τ , $\tau = 1, \dots, T$, $\lim_{\beta/\alpha \rightarrow 0} x(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \mathbf{\Gamma}, \mathbf{H}^{(\tau)})$ exists and equals to a function

$$\bar{x}_{\mathbf{\Gamma}}(\mathbf{H}^{(\tau)}\mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}), \mathbf{d}, \mathbf{Q}_{21}, \boldsymbol{\xi}). \quad (20)$$

Then

$$\begin{aligned} & \lim_{\beta/\alpha \rightarrow 0} E[x(\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta)] \\ &= E[\bar{x}_{\mathbf{\Gamma}}(\operatorname{diag}(\mathbf{G}_{11}(\mathbf{W}_{11}), \mathbf{G}_{22}(\mathbf{W}_{22})), (\mathbf{d}_1(\mathbf{W}_{11}), \mathbf{d}_2(\mathbf{W}_{22})), \mathbf{Z}_{21}, \boldsymbol{\xi})], \end{aligned} \quad (21)$$

where the expectation on the right side of (21) is taken with respect to the following mutually independent distributions

$$\begin{aligned} \mathbf{W}_{11} &\sim \mathbf{W}_m(n, \boldsymbol{\Xi}_1), \\ \mathbf{W}_{22} &\sim \mathbf{W}_{p-m}(n-m, \boldsymbol{\Xi}_2), \\ \mathbf{Z}_{21} &\sim \mathbf{N}_{(p-m) \times m}(\mathbf{0}, \mathbf{I}_{p-m} \otimes \mathbf{I}_m), \end{aligned} \quad (22)$$

and $\mathbf{G}_{ss}(\mathbf{W}_{ss})$, $\mathbf{d}_s(\mathbf{W}_{ss})$, $s = 1, 2$, are the components in the unique spectral decomposition of \mathbf{W}_{ss} for $s = 1, 2$;

$$\begin{aligned} \mathbf{W}_{11} &= \mathbf{G}_{11}\mathbf{D}_1\mathbf{G}'_{11}, \quad \mathbf{D}_1 = \operatorname{diag}(d_1, \dots, d_m), \quad \mathbf{d}_1 = (d_1, \dots, d_m), \\ \mathbf{W}_{22} &= \mathbf{G}_{22}\mathbf{D}_2\mathbf{G}'_{22}, \quad \mathbf{D}_2 = \operatorname{diag}(d_{m+1}, \dots, d_p), \quad \mathbf{d}_2 = (d_{m+1}, \dots, d_p). \end{aligned} \quad (23)$$

The proof is given in Appendix.

The following theorem on the asymptotic distributions is actually a corollary of Lemma 1. Let

$$\begin{aligned} \widetilde{\mathbf{W}}_{11} &= \widetilde{\mathbf{G}}_{11}\mathbf{D}_1\widetilde{\mathbf{G}}'_{11}, \\ \widetilde{\mathbf{W}}_{22} &= \widetilde{\mathbf{G}}_{22}\mathbf{D}_2\widetilde{\mathbf{G}}'_{22}, \\ \widetilde{\mathbf{Z}}_{21} &= \alpha^{1/2}\beta^{-1/2}\boldsymbol{\Xi}_2^{-1/2}\widetilde{\mathbf{G}}_{21}\mathbf{D}_1^{1/2}, \end{aligned}$$

where all the elements on the right-hand side are defined in Section 1.

Theorem 2 As $\beta/\alpha \rightarrow 0$,

$$\begin{aligned} \widetilde{\mathbf{W}}_{11} &\xrightarrow{d} \mathbf{W}_m(n, \boldsymbol{\Xi}_1), \\ \widetilde{\mathbf{W}}_{22} &\xrightarrow{d} \mathbf{W}_{p-m}(n-m, \boldsymbol{\Xi}_2), \\ \widetilde{\mathbf{Z}}_{21} &\xrightarrow{d} \mathbf{N}_{(p-m) \times m}(\mathbf{0}, \mathbf{I}_{p-m} \otimes \mathbf{I}_m) \end{aligned}$$

and $\widetilde{\mathbf{W}}_{11}$, $\widetilde{\mathbf{W}}_{22}$, $\widetilde{\mathbf{Z}}_{21}$ are asymptotically mutually independently distributed.

Proof. Let $\Theta_{11} : m \times m$ symmetric matrix, $\Theta_{22} : (p - m) \times (p - m)$ symmetric matrix and $\Theta_{21} : m \times (p - m)$ matrix. Consider the moment generating function

$$\begin{aligned} x(\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) &= \exp(\text{tr } \widetilde{\mathbf{W}}_{11} \Theta_{11} + \text{tr } \widetilde{\mathbf{W}}_{22} \Theta_{22} + \text{tr } \widetilde{\mathbf{Z}}_{21} \Theta_{21}) \\ &= \exp\left(\sum_{s=1}^2 \text{tr } \widetilde{\mathbf{W}}_{ss} \Theta_{ss} + \text{tr } \widetilde{\mathbf{Z}}_{21} \Theta_{21}\right). \end{aligned}$$

For $\mathbf{H}^{(\tau)} = \text{diag}(\mathbf{H}_1^{(\tau)}, \mathbf{H}_2^{(\tau)})$, $\mathbf{H}_1^{(\tau)} \in \mathcal{O}^+(m)$, $\mathbf{H}_2^{(\tau)} \in \mathcal{O}^+(p - m)$, we have

$$\begin{aligned} x(\boldsymbol{\Gamma} \mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}), \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) &= \exp\left\{\sum_{s=1}^2 \text{tr}(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}))_{ss} \mathbf{D}_s (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}))'_{ss} \Theta_{ss} \right. \\ &\quad \left. + \text{tr } \alpha^{1/2} \beta^{-1/2} \boldsymbol{\Xi}_2^{-1/2} (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}))_{21} \mathbf{D}_1^{1/2} \Theta_{21}\right\}. \end{aligned}$$

From (5)

$$(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}))_{21} = \mathbf{H}_2^{(\tau)} \mathbf{U}_{21},$$

hence from (8)

$$\alpha^{1/2} \beta^{-1/2} \boldsymbol{\Xi}_2^{-1/2} (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}))_{21} \mathbf{D}_1^{1/2} = \mathbf{Q}_{21}.$$

This leads to

$$x(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \boldsymbol{\Gamma}, \mathbf{H}^{(\tau)}) = \exp\left\{\sum_{s=1}^2 \text{tr}(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}))_{ss} \mathbf{D}_s (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}))'_{ss} \Theta_{ss} + \text{tr } \mathbf{Q}_{21} \Theta_{21}\right\},$$

with $\mathbf{u} = \mathbf{u}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)$. Therefore from (18)

$$\begin{aligned} &\lim_{\beta/\alpha \rightarrow 0} x(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \boldsymbol{\Gamma}, \mathbf{H}^{(\tau)}) \\ &= \exp\left\{\sum_{s=1}^2 \text{tr}(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))_{ss} \mathbf{D}_s (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))'_{ss} \Theta_{ss} + \text{tr } \mathbf{Q}_{21} \Theta_{21}\right\}. \end{aligned}$$

From Lemma 1,

$$\begin{aligned} &\lim_{\beta/\alpha \rightarrow 0} E[\exp(\text{tr } \widetilde{\mathbf{W}}_{11} \Theta_{11} + \text{tr } \widetilde{\mathbf{W}}_{22} \Theta_{22} + \text{tr } \widetilde{\mathbf{Z}}_{21} \Theta_{21})] \\ &= E[\exp\left\{\sum_{s=1}^2 \text{tr } \mathbf{G}_{ss}(\mathbf{W}_{ss}) \mathbf{D}_s (\mathbf{W}_{ss}) \mathbf{G}_{ss}(\mathbf{W}_{ss})' \Theta_{ss} + \text{tr } \mathbf{Z}_{21} \Theta_{21}\right\}] \\ &= E[\text{etr } \mathbf{W}_{11} \Theta_{11}] E[\text{etr } \mathbf{W}_{22} \Theta_{22}] E[\text{etr } \mathbf{Z}_{21} \Theta_{21}], \end{aligned}$$

where in the second and third equations the expectations are taken with respect to the distributions (22) in Lemma 1. ■

2.3 Multi-block Partition

In this section, we extend Theorem 2 into multi-block cases. We partition $(1, \dots, p)$ into k blocks;

$$\begin{array}{ll} \text{1st block} & (m_0 + 1, \dots, m_1), \\ \text{2nd block} & (m_1 + 1, \dots, m_2), \\ & \vdots \\ \text{kth block} & (m_{k-1} + 1, \dots, m_k), \end{array}$$

where

$$m_0 = 0 < m_1 < m_2 < \cdots < m_k = p.$$

Let $[i]$, $i = 1, \dots, p$, denote the block containing i , i.e.,

$$[i] = s, \quad \text{if } m_{s-1} + 1 \leq i \leq m_s.$$

We also use the notations $\bar{m}_s = m_s - m_{s-1}$, $s = 1, \dots, k$, for the block sizes.

Correspondingly to the above partition, we make the following partition of a $p \times p$ matrix $\mathbf{A} = (a_{ij})$;

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_{11} & \cdots & \mathbf{A}_{1k} \\ \vdots & \ddots & \vdots \\ \mathbf{A}_{k1} & \cdots & \mathbf{A}_{kk} \end{pmatrix}, \quad \mathbf{A}_{st} : \bar{m}_s \times \bar{m}_t \text{ matrix, } 1 \leq s, t \leq k.$$

For a diagonal matrix $\mathbf{A} = \text{diag}(a_1, \dots, a_p)$, we use the notation

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_1 & & \mathbf{0} \\ & \ddots & \\ \mathbf{0} & & \mathbf{A}_k \end{pmatrix}, \quad \mathbf{A}_s = \text{diag}(a_{m_{s-1}+1}, \dots, a_{m_s}), \quad s = 1, \dots, k.$$

Consider the following parametrization of $\mathbf{l}, \boldsymbol{\lambda}$

$$\begin{aligned} \lambda_i &= \xi_i \alpha_{[i]}, \quad 1 \leq i \leq p. \\ l_i &= d_i \alpha_{[i]}, \quad 1 \leq i \leq p, \end{aligned}$$

In this subsection we again consider that ξ_i 's are fixed. Now we define $\widetilde{\mathbf{W}}_{ss}, \widetilde{\mathbf{Z}}_{st}$, $1 \leq t < s \leq k$;

$$\begin{aligned} \widetilde{\mathbf{W}}_{ss} &= \widetilde{\mathbf{G}}_{ss} \mathbf{D}_s \widetilde{\mathbf{G}}'_{ss}, \\ \widetilde{\mathbf{Z}}_{st} &= \alpha_t^{1/2} \alpha_s^{-1/2} \boldsymbol{\Xi}_s^{-1/2} \widetilde{\mathbf{G}}_{st} \mathbf{D}_t^{1/2}, \end{aligned}$$

where notations of the right-hand side are defined in Section 1. The following theorem is the extension of Theorem 2.

Theorem 3 As $(\alpha_2/\alpha_1, \alpha_3/\alpha_2, \dots, \alpha_k/\alpha_{k-1}) \rightarrow \mathbf{0}$,

$$\begin{aligned} \widetilde{\mathbf{W}}_{ss} &\xrightarrow{d} \mathbf{W}_{\alpha_s}(n - m_{s-1}, \boldsymbol{\Xi}_s), \quad 1 \leq s \leq k, \\ \widetilde{\mathbf{Z}}_{st} &\xrightarrow{d} \mathbf{N}_{\bar{m}_s \times \bar{m}_t}(\mathbf{0}, \mathbf{I}_{\bar{m}_s} \otimes \mathbf{I}_{\bar{m}_t}), \quad 1 \leq t < s \leq k, \end{aligned}$$

and $\widetilde{\mathbf{W}}_{ss} (1 \leq s \leq k), \widetilde{\mathbf{Z}}_{st} (1 \leq t < s \leq k)$ are asymptotically mutually independently distributed.

Proof. Though we can prove the theorem in the same manner as the proof of Theorem 2, it is notationally too cumbersome. Instead we will prove the theorem by using Theorem 2 recursively. Let $r_1 = \alpha_1$ and $r_t = \alpha_t/\alpha_{t-1}$, $t = 2, \dots, k$, then $\prod_{t=1}^s r_t = \alpha_s$, $s = 1, \dots, k$. Note for $1 \leq i \leq p$,

$$l_i = d_i \alpha_{[i]} = d_i \prod_{t=1}^{[i]} r_t, \quad \lambda_i = \xi_i \alpha_{[i]} = \xi_i \prod_{t=1}^{[i]} r_t.$$

We consider the moment generating function

$$E\left[\exp\left(\text{tr} \sum_{s=1}^k \widetilde{\mathbf{W}}_{ss} \boldsymbol{\Theta}_{ss} + \text{tr} \sum_{1 \leq t < s \leq k} \widetilde{\mathbf{Z}}_{st} \boldsymbol{\Theta}_{st}\right)\right],$$

where $\boldsymbol{\Theta}_{ss} (1 \leq s \leq k)$ and $\boldsymbol{\Theta}_{st} (1 \leq t < s \leq k)$ are respectively a $\bar{m}_s \times \bar{m}_s$ symmetric matrix and a $\bar{m}_t \times \bar{m}_s$ matrix. We have

$$\begin{aligned} & \lim_{(a_2/a_1, \dots, a_k/a_{k-1}) \rightarrow 0} E\left[\exp\left(\text{tr} \sum_{s=1}^k \widetilde{\mathbf{W}}_{ss} \boldsymbol{\Theta}_{ss} + \text{tr} \sum_{1 \leq t < s \leq k} \widetilde{\mathbf{Z}}_{st} \boldsymbol{\Theta}_{st}\right)\right] \\ &= \lim_{(r_2, \dots, r_k) \rightarrow 0} E\left[\exp\left(\text{tr} \sum_{s=1}^k \widetilde{\mathbf{W}}_{ss} \boldsymbol{\Theta}_{ss} + \text{tr} \sum_{1 \leq t < s \leq k} \widetilde{\mathbf{Z}}_{st} \boldsymbol{\Theta}_{st}\right)\right] \\ &= \lim_{r_2 \rightarrow 0} \cdots \lim_{r_k \rightarrow 0} E\left[\exp\left(\text{tr} \sum_{s=1}^k \widetilde{\mathbf{W}}_{ss} \boldsymbol{\Theta}_{ss} + \text{tr} \sum_{1 \leq t < s \leq k} \widetilde{\mathbf{Z}}_{st} \boldsymbol{\Theta}_{st}\right)\right] \end{aligned}$$

We omit technical arguments on uniform convergences, which guarantees the decomposition of $\lim_{(r_2, \dots, r_k) \rightarrow 0}$ in the second line into step by step limiting operations $\lim_{r_2 \rightarrow 0} \cdots \lim_{r_k \rightarrow 0}$ in the third line.

Consider the partitions;

$$\widetilde{\mathbf{G}} = \begin{pmatrix} & & & \widetilde{\mathbf{G}}_{1k} \\ & \widetilde{\mathbf{G}}^{(k-1)} & & \vdots \\ & & & \widetilde{\mathbf{G}}_{k-1k} \\ \widetilde{\mathbf{G}}_{k1} & \cdots & \widetilde{\mathbf{G}}_{kk-1} & \widetilde{\mathbf{G}}_{kk} \end{pmatrix} \quad \text{where} \quad \widetilde{\mathbf{G}}^{(k-1)} = \begin{pmatrix} \widetilde{\mathbf{G}}_{11} & \cdots & \widetilde{\mathbf{G}}_{1k-1} \\ \vdots & \ddots & \vdots \\ \widetilde{\mathbf{G}}_{k-11} & \cdots & \widetilde{\mathbf{G}}_{k-1k-1} \end{pmatrix}.$$

Define \mathbf{D}^* , $\boldsymbol{\Xi}^*$ as partitioned matrices;

$$\mathbf{D}^* = \begin{pmatrix} \mathbf{L}^{(k-1)} & \mathbf{0} \\ \mathbf{0} & \mathbf{D}_k \end{pmatrix}, \quad \boldsymbol{\Xi}^* = \begin{pmatrix} \boldsymbol{\Lambda}^{(k-1)} & \mathbf{0} \\ \mathbf{0} & \boldsymbol{\Xi}_k \end{pmatrix},$$

where

$$\mathbf{L}^{(k-1)} = \text{diag}(l_1, \dots, l_{m_{k-1}}), \quad \boldsymbol{\Lambda}^{(k-1)} = \text{diag}(\lambda_1, \dots, \lambda_{m_{k-1}}).$$

Let $\alpha = 1$, $\beta = \alpha_k = \prod_{t=1}^k r_t$. Then

$$\mathbf{L} = \begin{pmatrix} \mathbf{L}^{(k-1)}\alpha & \mathbf{0} \\ \mathbf{0} & \mathbf{D}_k\beta \end{pmatrix}, \quad \boldsymbol{\Lambda} = \begin{pmatrix} \boldsymbol{\Lambda}^{(k-1)}\alpha & \mathbf{0} \\ \mathbf{0} & \boldsymbol{\Xi}_k\beta \end{pmatrix}.$$

Since as $r_k \rightarrow 0$, $\beta/\alpha \rightarrow 0$, from Theorem 2, we have

$$\begin{aligned} \mathbf{S}^{(k-1)} &= \widetilde{\mathbf{G}}^{(k-1)} \mathbf{L}^{(k-1)} \widetilde{\mathbf{G}}^{(k-1)'} \xrightarrow{d} \mathbf{W}_{m_{k-1}}(n, \boldsymbol{\Lambda}^{(k-1)}), \\ \widetilde{\mathbf{W}}_{kk} &\xrightarrow{d} \mathbf{W}_{\bar{m}_k}(n - m_{k-1}, \boldsymbol{\Xi}_k), \\ \widetilde{\mathbf{Z}}_{kt} &\xrightarrow{d} \mathbf{N}_{\bar{m}_k \times \bar{m}_t}(\mathbf{0}, \mathbf{I}_{\bar{m}_k} \otimes \mathbf{I}_{\bar{m}_t}), \quad 1 \leq t \leq k-1, \end{aligned}$$

and the asymptotic distributions are mutually independent. Therefore

$$\begin{aligned} & \lim_{r_k \rightarrow 0} E \left[\exp \left(\text{tr} \sum_{s=1}^k \widetilde{\mathbf{W}}_{ss} \boldsymbol{\Theta}_{ss} + \text{tr} \sum_{1 \leq t < s \leq k} \widetilde{\mathbf{Z}}_{st} \boldsymbol{\Theta}_{st} \right) \right] \\ &= E \left[\exp \left(\text{tr} \sum_{s=1}^{k-1} \widetilde{\mathbf{W}}_{ss} (\mathbf{S}^{(k-1)}) \boldsymbol{\Theta}_{ss} + \text{tr} \sum_{1 \leq t < s \leq k-1} \widetilde{\mathbf{Z}}_{st} (\mathbf{S}^{(k-1)}) \boldsymbol{\Theta}_{st} \right) \right] \\ & \quad \times E \left[\text{etr} \widetilde{\mathbf{W}}_{kk} \boldsymbol{\Theta}_{kk} \right] \times \prod_{t=1}^{k-1} E \left[\text{etr} \widetilde{\mathbf{Z}}_{kt} \boldsymbol{\Theta}_{kt} \right], \end{aligned}$$

where the expectations on the right-hand side is taken with respect to the above asymptotic distributions. If we apply Theorem 2 again to $\mathbf{S}^{(k-1)}$ and recursively to the upper-left block Wishart distribution which asymptotically arises, we gain the result. \blacksquare

Note that Theorem 3 reduces to Theorem 2 of Takemura and Sheena (2005) for the extreme case of 1-element blocks $\bar{m}_s = 1$, $s = 1, \dots, p$. Therefore Theorem 3 is a generalization of Theorem 2 of Takemura and Sheena (2005).

3 Application to Estimation of $\boldsymbol{\Sigma}$

3.1 Loss Functions and Orthogonally Equivariant Estimators

In this section, we apply the asymptotic result on the distribution of \mathbf{S} to the estimation of $\boldsymbol{\Sigma}$ when β/α vanishes. We take a decision-theoretic approach to evaluate the performance of the estimators. We deal with the two loss functions; one is Stein's loss (entropy loss) function

$$L_1(\widehat{\boldsymbol{\Sigma}}, \boldsymbol{\Sigma}) = \text{tr}(\widehat{\boldsymbol{\Sigma}}\boldsymbol{\Sigma}^{-1}) - \log |\widehat{\boldsymbol{\Sigma}}\boldsymbol{\Sigma}^{-1}| - p, \quad (24)$$

and the other is a scale-invariant quadratic loss function

$$L_2(\widehat{\boldsymbol{\Sigma}}, \boldsymbol{\Sigma}) = \text{tr}(\widehat{\boldsymbol{\Sigma}}\boldsymbol{\Sigma}^{-1} - \mathbf{I}_p)^2. \quad (25)$$

The associated risk functions are denoted as

$$R_d(\widehat{\boldsymbol{\Sigma}}, \boldsymbol{\Sigma}) = E[L_d(\widehat{\boldsymbol{\Sigma}}, \boldsymbol{\Sigma})], \quad d = 1, 2.$$

The classical estimator of $\boldsymbol{\Sigma}$ is the unbiased estimator

$$\widehat{\boldsymbol{\Sigma}}^U = n^{-1} \mathbf{S},$$

which has been widely used for many statistical analysis, especially with statistical software packages. However, as James and Stein (1961) showed, this estimator is neither minimax nor admissible with Stein's loss function (24). The same drawback with respect to the quadratic loss function (25) was reported by Olkin and Selliah (1977). Following these initiative papers, much literature has been written seeking for a superior estimator to $\widehat{\boldsymbol{\Sigma}}^U$. See Pal (1993) for the review on the estimation of $\boldsymbol{\Sigma}$. In this paper we only refer to orthogonally equivariant estimators proposed by

Stein (1977), Dey and Srinivasan (1985) and Krishnamoorthy and Gupta (1989). An estimator of the form

$$\widehat{\Sigma} = \mathbf{G}\Psi(\mathbf{L})\mathbf{G}', \quad \Psi(\mathbf{L}) = \text{diag}(\psi_1(\mathbf{l}), \dots, \psi_p(\mathbf{l}))$$

is called *orthogonally equivariant*; i.e., $\widehat{\Sigma}(\mathbf{G}\mathbf{S}\mathbf{G}') = \mathbf{G}\widehat{\Sigma}(\mathbf{S})\mathbf{G}'$, $\forall \mathbf{G} \in \mathcal{O}(p)$.

Stein (1977) and Dey and Srinivasan (1985) proposed the orthogonally equivariant estimator, $\widehat{\Sigma}^{SDS}$, defined by

$$\psi_i(\mathbf{l}) = l_i \Delta_i^{JS}, \quad 1 \leq i \leq p,$$

where $\Delta_i^{JS} = (n+p+1-2i)^{-1}$. $\widehat{\Sigma}^{SDS}$ is of simple form but dominates $\widehat{\Sigma}^U$ with substantially better risk w.r.t the loss function (24). It is also a minimax estimator. See Dey and Srinivasan (1985) and Sugiura and Ishibayashi (1997) for more details. Order preservation among $\psi_i(\mathbf{l})$, $i = 1, \dots, p$, is discussed in Sheena and Takemura (1992).

The orthogonally equivariant estimator $\widehat{\Sigma}^{KG}$ is defined by

$$\psi_i(\mathbf{l}) = l_i \Delta_i^{OS}, \quad 1 \leq i \leq p,$$

where Δ_i^{OS} is given by

$$(\Delta_1^{OS}, \dots, \Delta_p^{OS})' = \mathbf{A}^{-1}\mathbf{b}$$

with a $p \times p$ matrix $\mathbf{A} = (a_{ij})$ and a $p \times 1$ vector $\mathbf{b} = (b_i)$ defined by

$$a_{ij} = \begin{cases} (n+p-2i+1)(n+p-2i+3), & \text{if } i=j, \\ (n+p-2i+1), & \text{if } i>j, \\ (n+p-2j+1), & \text{if } j>i, \end{cases}$$

$$b_i = n+p+1-2i, \quad i = 1, \dots, p.$$

$\widehat{\Sigma}^{KG}$ is conjectured to be a minimax estimator which dominates $\widehat{\Sigma}^U$ w.r.t. the loss function (25). This was proved by Sheena (2002) for the case $p = 2$.

In this section we only consider orthogonally equivariant estimators given by

$$\psi_i(\mathbf{l}) = c_i l_i, \quad 1 \leq i \leq p \tag{26}$$

with some constant c_i ($1 \leq i \leq p$), or in the matrix expression,

$$\Psi(\mathbf{L}) = \mathbf{L}^{1/2} \mathbf{C} \mathbf{L}^{1/2}, \quad \mathbf{C} = \text{diag}(c_1, \dots, c_p).$$

It is interesting that $\widehat{\Sigma}^{SDS}$ and $\widehat{\Sigma}^{KG}$ are also the minimum risk estimators among the estimators of the form (26) respectively for $L_1(\cdot, \cdot)$ and $L_2(\cdot, \cdot)$ when all the population eigenvalues are dispersed. See Takemura and Sheena (2005) for more details.

3.2 Asymptotic Risk

This subsection is devoted to the calculation of the asymptotic risks $\widetilde{R}_d(\widehat{\Sigma}, \Sigma)$

$$\widetilde{R}_d(\widehat{\Sigma}, \Sigma) = \lim_{\beta/\alpha \rightarrow 0} R_d(\widehat{\Sigma}, \Sigma), \quad d = 1, 2,$$

for an orthogonally equivariant estimator defined by (26). Note that

$$\begin{aligned} R_1(\widehat{\Sigma}, \Sigma) &= E[\text{tr } \mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \Gamma \Lambda^{-1} \Gamma'] - \log |\mathbf{C}| - E[\log |\Sigma^{-1/2} \mathbf{S} \Sigma^{-1/2}|] - p \\ &= E[\text{tr } \mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \Gamma \Lambda^{-1} \Gamma'] - \sum_{i=1}^p \log c_i - \sum_{i=1}^p E[\log \chi_{n-i+1}^2] - p. \end{aligned} \quad (27)$$

$$\begin{aligned} R_2(\widehat{\Sigma}, \Sigma) &= E[\text{tr}(\mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \Gamma \Lambda^{-1} \Gamma' - \mathbf{I}_p)^2] \\ &= E[\text{tr}(\mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \Gamma \Lambda^{-1} \Gamma')^2] - 2E[\text{tr } \mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \Gamma \Lambda^{-1} \Gamma'] + p. \end{aligned} \quad (28)$$

For the evaluation $E[\log |\Sigma^{-1/2} \mathbf{S} \Sigma^{-1/2}|]$, see e.g. (10) in p.132 of Muirhead (1982).

We start with the following lemma, the proof of which is given in Appendix.

Lemma 2

$$\begin{aligned} &\lim_{\beta/\alpha \rightarrow 0} E[\text{tr } \mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \Gamma \Lambda^{-1} \Gamma'] \\ &= E[\text{tr } \mathbf{G}_{11} \mathbf{D}_1^{1/2} \mathbf{C}_1 \mathbf{D}_1^{1/2} \mathbf{G}'_{11} \Xi_1^{-1}] + E[\text{tr } \mathbf{G}_{22} \mathbf{D}_2^{1/2} \mathbf{C}_2 \mathbf{D}_2^{1/2} \mathbf{G}'_{22} \Xi_2^{-1}] \\ &\quad + (p - m) \text{tr } \mathbf{C}_1, \end{aligned} \quad (29)$$

$$\begin{aligned} &\lim_{\beta/\alpha \rightarrow 0} E[\text{tr}(\mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \Gamma \Lambda^{-1} \Gamma')^2] \\ &= E[\text{tr}(\mathbf{G}_{11} \mathbf{D}_1^{1/2} \mathbf{C}_1 \mathbf{D}_1^{1/2} \mathbf{G}'_{11} \Xi_1^{-1})^2] + E[\text{tr}(\mathbf{G}_{22} \mathbf{D}_2^{1/2} \mathbf{C}_2 \mathbf{D}_2^{1/2} \mathbf{G}'_{22} \Xi_2^{-1})^2] \\ &\quad + 2(p - m) E[\text{tr } \mathbf{C}_1^2 \mathbf{D}_1^{1/2} \mathbf{G}'_{11} \Xi_1^{-1} \mathbf{G}_{11} \mathbf{D}_1^{1/2}] + 2 \text{tr } \mathbf{C}_1 E[\text{tr } \Xi_2^{-1} \mathbf{G}_{22} \mathbf{D}_2^{1/2} \mathbf{C}_2 \mathbf{D}_2^{1/2} \mathbf{G}'_{22}] \\ &\quad + (p - m)(p - m + 2) \sum_{i=1}^m c_i^2 + 2(p - m) \sum_{1 \leq i < s \leq m} c_i c_s, \end{aligned} \quad (30)$$

where the expectations on the right-hand side in (29) and (30) are taken with respect to the distributions in (22) and the decompositions in (23).

Now suppose that under the distribution of \mathbf{W}_{ss} , $s = 1, 2$, in (22) and their spectral decomposition in (23), we estimate Ξ_s , $s = 1, 2$, by the following orthogonally equivariant estimators

$$\begin{aligned} \widehat{\Xi}_1 &= \mathbf{G}_{11} \mathbf{D}_1^{1/2} \mathbf{C}_1 \mathbf{D}_1^{1/2} \mathbf{G}'_{11}, & \mathbf{C}_1 &= \text{diag}(c_1, \dots, c_m), \\ \widehat{\Xi}_2 &= \mathbf{G}_{22} \mathbf{D}_2^{1/2} \mathbf{C}_2 \mathbf{D}_2^{1/2} \mathbf{G}'_{22}, & \mathbf{C}_2 &= \text{diag}(c_{m+1}, \dots, c_p), \end{aligned}$$

then the risks w.r.t. each loss function (24), (25) are given by

$$\begin{aligned} R_{11}(\widehat{\Xi}_1, \Xi_1) &= E[\text{tr}(\widehat{\Xi}_1 \Xi_1^{-1}) - \log |\widehat{\Xi}_1 \Xi_1^{-1}| - m], \\ R_{21}(\widehat{\Xi}_2, \Xi_2) &= E[\text{tr}(\widehat{\Xi}_2 \Xi_2^{-1}) - \log |\widehat{\Xi}_2 \Xi_2^{-1}| - p + m], \\ R_{12}(\widehat{\Xi}_1, \Xi_1) &= E[\text{tr}(\widehat{\Xi}_1 \Xi_1^{-1} - \mathbf{I}_m)^2], \\ R_{22}(\widehat{\Xi}_2, \Xi_2) &= E[\text{tr}(\widehat{\Xi}_2 \Xi_2^{-1} - \mathbf{I}_{p-m})^2]. \end{aligned}$$

The following theorem gives the decomposition of the asymptotic risk, $\widetilde{R}_d(\widehat{\Sigma}, \Sigma)$, into the risks R_{1d} , R_{2d} and the residuals R_{3d} for $d = 1, 2$.

Theorem 4 For $d = 1, 2$,

$$\tilde{R}_d(\hat{\Sigma}, \Sigma) = R_{1d}(\hat{\Xi}_1, \Xi_1) + R_{2d}(\hat{\Xi}_2, \Xi_2) + R_{3d},$$

where

$$R_{31} = (p - m) \sum_{i=1}^m c_i,$$

and

$$\begin{aligned} R_{32} &= 2(p - m)E[\text{tr } \mathbf{C}_1^2 \mathbf{D}_1^{1/2} \mathbf{G}'_{11} \Xi_1^{-1} \mathbf{G}_{11} \mathbf{D}_1^{1/2}] + 2 \text{tr } \mathbf{C}_1 E[\text{tr } \Xi_2^{-1} \mathbf{G}_{22} \mathbf{D}_2^{1/2} \mathbf{C}_2 \mathbf{D}_2^{1/2} \mathbf{G}'_{22}] \\ &\quad + (p - m)(p - m + 2) \sum_{i=1}^m c_i^2 + 2(p - m) \sum_{1 \leq i < s \leq m} c_i c_s - 2(p - m) \sum_{i=1}^m c_i. \end{aligned}$$

All the expectations are taken with respect to the distributions (22) and the decompositions (23).

Proof. From (27),

$$\begin{aligned} R_{11}(\hat{\Sigma}_1, \Sigma_1) &= E[\text{tr } \mathbf{G}_{11} \mathbf{D}_1^{1/2} \mathbf{C}_1 \mathbf{D}_1^{1/2} \mathbf{G}'_{11} \Xi_1^{-1}] - \sum_{i=1}^m \log c_i - \sum_{i=1}^m E[\log \chi_{n-i+1}^2] - m, \\ R_{21}(\hat{\Sigma}_2, \Sigma_2) &= E[\text{tr } \mathbf{G}_{22} \mathbf{D}_2^{1/2} \mathbf{C}_2 \mathbf{D}_2^{1/2} \mathbf{G}'_{22} \Xi_2^{-1}] - \sum_{i=m+1}^p \log c_i - \sum_{i=m+1}^p E[\log \chi_{n-i+1}^2] - p + m. \end{aligned}$$

Using (29) together with the above result, we have the result for $\tilde{R}_1(\hat{\Sigma}, \Sigma)$. From (28),

$$\begin{aligned} R_{12}(\hat{\Sigma}_1, \Sigma_1) &= E[\text{tr}(\mathbf{G}_{11} \mathbf{D}_1^{1/2} \mathbf{C}_1 \mathbf{D}_1^{1/2} \mathbf{G}'_{11} \Xi_1^{-1})^2] - 2E[\text{tr } \mathbf{G}_{11} \mathbf{D}_1^{1/2} \mathbf{C}_1 \mathbf{D}_1^{1/2} \mathbf{G}'_{11} \Xi_1^{-1}] + m, \\ R_{22}(\hat{\Sigma}_2, \Sigma_2) &= E[\text{tr}(\mathbf{G}_{22} \mathbf{D}_2^{1/2} \mathbf{C}_2 \mathbf{D}_2^{1/2} \mathbf{G}'_{22} \Xi_2^{-1})^2] - 2E[\text{tr } \mathbf{G}_{22} \mathbf{D}_2^{1/2} \mathbf{C}_2 \mathbf{D}_2^{1/2} \mathbf{G}'_{22} \Xi_2^{-1}] + p - m. \end{aligned}$$

Using (30) and (29) together with the above result, we have the result for $\tilde{R}_2(\hat{\Sigma}, \Sigma)$. ■

3.3 Minimum Asymptotic Risk Estimator

Consider the model (1) and suppose $\tau_1 = \dots = \tau_m (= \tau)$ in (2). Then $\alpha = \tau + \sigma^2$ and $\beta = \sigma^2$ and

$$\Xi_1 = \mathbf{I}_m, \quad \Xi_2 = \mathbf{I}_{p-m}. \quad (31)$$

This assumption may not be very realistic. However note that it is trivially satisfied in the one-factor model $m = 1$, which is frequently used in practice. In this subsection we focus on the estimation of Σ under the condition (31). In this case, since we have no unknown parameters anymore, the asymptotic risk is uniquely determined, hence we can derive the “best” i.e., minimum asymptotic risk estimator among the orthogonally equivariant estimators of the form (26). The following theorem gives the asymptotic risk for the case (31).

Theorem 5 If $\Xi_1 = \mathbf{I}_m$, $\Xi_2 = \mathbf{I}_{p-m}$, then the asymptotic risk $\tilde{R}_d(\hat{\Sigma}, \Sigma)$, $d = 1, 2$, is given by the following function of $\mathbf{c} = (c_1, \dots, c_p)'$.

$$\tilde{R}_1(\hat{\Sigma}, \Sigma) = \sum_{i=1}^p (b_i c_i - \log c_i) - \sum_{i=1}^p E[\log \chi_{n-i+1}^2] - p, \quad (32)$$

$$\tilde{R}_2(\hat{\Sigma}, \Sigma) = \mathbf{c}' \mathbf{A} \mathbf{c} - 2\mathbf{b}' \mathbf{c} + p, \quad (33)$$

where $\mathbf{b} = (b_1, \dots, b_p)'$ is given by

$$b_i = \begin{cases} E[d_i] + p - m, & \text{if } 1 \leq i \leq m, \\ E[d_i], & \text{if } m + 1 \leq i \leq p, \end{cases}$$

and $p \times p$ symmetric matrix $\mathbf{A} = (a_{ij})$ is given by

$$a_{ij} = \begin{cases} E[d_i^2 + 2(p - m)d_i] + (p - m)(p - m + 2), & \text{if } 1 \leq i = j \leq m, \\ E[d_i^2], & \text{if } m + 1 \leq i = j \leq p, \\ p - m, & \text{if } 1 \leq i \neq j \leq m, \\ E[d_j], & \text{if } 1 \leq i \leq m < j \leq p, \\ E[d_i], & \text{if } 1 \leq j \leq m < i \leq p, \\ 0, & \text{otherwise.} \end{cases}$$

All the expectations are taken with respect to the distribution (22) and the decompositions (23) with $\mathbf{\Xi}_1 = \mathbf{I}_m$, $\mathbf{\Xi}_2 = \mathbf{I}_{p-m}$.

Proof. Evaluating $R_{jd}(\hat{\Sigma}_j, \Sigma_j)$, $1 \leq j, d \leq 2$ in Theorem 4 when $\mathbf{\Xi}_1 = \mathbf{I}_m$, $\mathbf{\Xi}_2 = \mathbf{I}_{p-m}$, we have the following results.

$$\begin{aligned} R_{11}(\hat{\mathbf{\Xi}}_1, \mathbf{\Xi}_1) &= E[L_1(\hat{\mathbf{\Xi}}_1, \mathbf{I}_m)] = E[\text{tr} \hat{\mathbf{\Xi}}_1 - \log |\hat{\mathbf{\Xi}}_1| - m] \\ &= E\left[\sum_{i=1}^m d_i c_i - \log |\mathbf{W}_{11}|\right] - \sum_{i=1}^m \log c_i - m \\ &= \sum_{i=1}^m E[d_i] c_i - E[\log |\mathbf{W}_{11}|] - \sum_{i=1}^m \log c_i - m. \\ R_{21}(\hat{\mathbf{\Xi}}_2, \mathbf{\Xi}_2) &= \sum_{i=m+1}^p E[d_i] c_i - E[\log |\mathbf{W}_{22}|] - \sum_{i=m+1}^p \log c_i - p + m. \\ R_{12}(\hat{\mathbf{\Xi}}_1, \mathbf{\Xi}_1) &= E[L_2(\hat{\mathbf{\Xi}}_1, \mathbf{I}_m)] = E[\text{tr}(\hat{\mathbf{\Xi}}_1 - \mathbf{I}_m)^2] \\ &= E[\text{tr} \hat{\mathbf{\Xi}}_1^2 - 2 \text{tr} \hat{\mathbf{\Xi}}_1] + m = E\left[\sum_{i=1}^m d_i^2 c_i^2 - 2 \sum_{i=1}^m d_i c_i\right] + m \\ &= \sum_{i=1}^m E[d_i^2] c_i^2 - 2 \sum_{i=1}^m E[d_i] c_i + m. \\ R_{22}(\hat{\mathbf{\Xi}}_2, \mathbf{\Xi}_2) &= \sum_{i=m+1}^p E[d_i^2] c_i^2 - 2 \sum_{i=m+1}^p E[d_i] c_i + p - m. \end{aligned}$$

Next we calculate R_{32} in Theorem 4 when $\mathbf{\Xi}_1 = \mathbf{I}_m$, $\mathbf{\Xi}_2 = \mathbf{I}_{p-m}$. Note that

$$\begin{aligned} 2(p - m)E[\text{tr} \mathbf{C}_1^2 \mathbf{D}_1^{1/2} \mathbf{G}'_{11} \mathbf{\Xi}_1^{-1} \mathbf{G}_{11} \mathbf{D}_1^{1/2}] &= 2(p - m)E[\text{tr} \mathbf{C}_1^2 \mathbf{D}_1] \\ &= 2(p - m) \sum_{i=1}^m E[d_i] c_i^2, \\ 2 \text{tr} \mathbf{C}_1 E[\text{tr} \mathbf{\Xi}_2^{-1} \mathbf{G}_{22} \mathbf{D}_2^{1/2} \mathbf{C}_2 \mathbf{D}_2^{1/2} \mathbf{G}'_{22}] &= 2 \left(\sum_{i=1}^m c_i \right) \left(\sum_{i=m+1}^p E[d_i] c_i \right). \end{aligned}$$

Therefore

$$R_{32} = \sum_{i=1}^m c_i^2 \{(p-m)(p-m+2) + 2(p-m)E[d_i]\} \\ + 2(p-m) \sum_{1 \leq i < s \leq m} c_i c_s + 2 \sum_{1 \leq i \leq m < s \leq p} c_i c_s E[d_s] - 2(p-m) \sum_{i=1}^m c_i.$$

Combining above results, we see that (32) and (33) hold. \blacksquare

Corollary 1 *The minimum asymptotic risk with respect to the loss function $L_1(\cdot, \cdot)$ is given by*

$$\sum_{i=1}^p \log b_i - \sum_{i=1}^p E[\log \chi_{n-i-1}^2].$$

It is attained by $\widehat{\Sigma}^{MA_1}$ given by $c_i = b_i^{-1}$, $i = 1, \dots, p$. The minimum asymptotic risk with respect to the loss function $L_2(\cdot, \cdot)$ is given by

$$p - \mathbf{b}' \mathbf{A}^{-1} \mathbf{b}.$$

It is attained by $\widehat{\Sigma}^{MA_2}$ given by $\mathbf{c} = \mathbf{A}^{-1} \mathbf{b}$.

Proof. The results are easily obtained by the minimization $\sum_{i=1}^p (b_i c_i - \log c_i)$ or $\mathbf{c}' \mathbf{A} \mathbf{c} - 2\mathbf{b}' \mathbf{c}$. \blacksquare

The calculation of the asymptotic risks in Theorem 5 and the c_i 's of $\widehat{\Sigma}^{MA_1}$ and $\widehat{\Sigma}^{MA_2}$ requires the evaluation of $E[d_i]$, $E[d_i^2]$, $i = 1, \dots, p$, that is, the first and the second moment of the eigenvalues of the Wishart distribution with the identity covariance matrix. Generally we need to make use of Monte Carlo simulation or numerical integration for the evaluation of the moments of the eigenvalues. However when p is small and n is appropriately even or odd depending on p , the analytic evaluation is feasible. See Section A.3 in Appendix for this evaluation.

Tables 1–5 give c_i 's for $\widehat{\Sigma}^U$, $\widehat{\Sigma}^{SDS}$, $\widehat{\Sigma}^{KG}$, $\widehat{\Sigma}^{MA_1}$, $\widehat{\Sigma}^{MA_2}$ when $p = 3, 4$ with several values of n . The value of c_i 's for the minimum asymptotic risk estimators $\widehat{\Sigma}^{MA_1}$, $\widehat{\Sigma}^{MA_2}$ is calculated by the aforementioned analytic method. Note that for the case $p = 2$, the minimum asymptotic risk estimator naturally coincides with $\widehat{\Sigma}^{SDS}$ ($\widehat{\Sigma}^{KG}$) which is the minimum asymptotic risk estimator for $L_1(L_2)$ when we see the total dispersion of population eigenvalues (see Takemura and Sheena (2005)). As it is well known, $n^{-1}l_i$ ($i = 1, \dots, p$) tends to overestimate the corresponding eigenvalue of Σ when i is small, while it tends to underestimate the corresponding eigenvalue of Σ when i is large. The estimators $\widehat{\Sigma}^{SDS}$, $\widehat{\Sigma}^{KG}$ modify this tendency by increasing weight $c_1 < \dots < c_p$. It is seen from the tables that $\widehat{\Sigma}^{MA_1}$, $\widehat{\Sigma}^{MA_2}$ enlarge the weight difference within each block in most cases; for example when $p = 4, m = 2$, the relation between c_i 's of $\widehat{\Sigma}^{SDS}$ ($\widehat{\Sigma}^{KG}$) (say c_i^{SDS} (c_i^{KG}), $i = 1, \dots, 4$) and those of $\widehat{\Sigma}^{MA_1}$ ($\widehat{\Sigma}^{MA_2}$) (say $c_i^{MA_1}$ ($c_i^{MA_2}$), $i = 1, \dots, 4$) is found as

$$c_1^{MA_1} < c_1^{SDS} < c_2^{SDS} < c_2^{MA_1}, \quad c_3^{MA_1} < c_3^{SDS} < c_4^{SDS} < c_4^{MA_1},$$

and

$$c_1^{MA_2} < c_1^{KG} < c_2^{KG} < c_2^{MA_2}, \quad c_3^{MA_2} < c_3^{KG} < c_4^{KG} < c_4^{MA_2}.$$

The tables also give asymptotic risk comparison w.r.t. L_1 among the estimators $\widehat{\Sigma}^U$, $\widehat{\Sigma}^{SDS}$, $\widehat{\Sigma}^{MA_1}$ (see ‘‘Asy.Risk1’’) and that w.r.t. L_2 among the estimators $\widehat{\Sigma}^U$, $\widehat{\Sigma}^{KG}$, $\widehat{\Sigma}^{MA_2}$ (see ‘‘Asy.Risk2’’).

Table 1: $p = 3, m = 1$

$n = 4$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 6$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.2500	0.1667	0.1060	0.1667	0.1019	c_1	0.1667	0.1250	0.0855	0.1250	0.0828
c_2	0.2500	0.2500	0.1332	0.2000	0.1321	c_2	0.1667	0.1667	0.1030	0.1304	0.0977
c_3	0.2500	0.5000	0.1902	1.0000	0.4491	c_3	0.1667	0.2500	0.1352	0.4286	0.2675
Asy.Risk1	2.1969	1.6592		1.4392		Asy.Risk1	1.2387	0.9820		0.8270	
R.R.R.		24.47		34.49		R.R.R.		20.72		33.23	
Asy.Risk2	3.0000		1.4120		1.2792	Asy.Risk2	2.0000		1.1056		0.9644
R.R.R.			52.93		57.36	R.R.R.			44.72		51.78

$n = 8$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 10$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.1250	0.1000	0.0724	0.1000	0.0707	c_1	0.1000	0.0833	0.0630	0.0833	0.0619
c_2	0.1250	0.1250	0.0849	0.0980	0.0782	c_2	0.1000	0.1000	0.0723	0.0790	0.0655
c_3	0.1250	0.1667	0.1053	0.2632	0.1878	c_3	0.1000	0.1250	0.0865	0.1872	0.1438
Asy.Risk1	0.8749	0.7187		0.5966		Asy.Risk1	0.6765	0.5692		0.4676	
R.R.R.		17.85		31.81		R.R.R.		15.85		30.88	
Asy.Risk2	1.5000		0.9140		0.7812	Asy.Risk2	1.2000		0.7817		0.6591
R.R.R.			39.07		47.92	R.R.R.			34.86		45.07

$n = 20$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 50$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.0500	0.0455	0.0385	0.0455	0.0383	c_1	0.0200	0.0192	0.0179	0.0192	0.0178
c_2	0.0500	0.0500	0.0418	0.0410	0.0370	c_2	0.0200	0.0200	0.0185	0.0173	0.0166
c_3	0.0500	0.0556	0.0460	0.0735	0.0647	c_3	0.0200	0.0208	0.0193	0.0248	0.0236
Asy.Risk1	0.3164	0.2819		0.2251		Asy.Risk1	0.1236	0.1155		0.0901	
R.R.R.		10.89		28.84		R.R.R.		6.51		27.05	
Asy.Risk2	0.6000		0.4598		0.3745	Asy.Risk2	0.2400		0.2093		0.1647
R.R.R.			23.37		37.59	R.R.R.			12.79		31.39

Table 2: $p = 3, m = 2$

$n = 5$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 7$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.2000	0.1429	0.0944	0.1154	0.0891	c_1	0.1429	0.1111	0.0784	0.0893	0.0726
c_2	0.2000	0.2000	0.1158	0.3000	0.1791	c_2	0.1429	0.1429	0.0930	0.2083	0.1417
c_3	0.2000	0.3333	0.1580	0.3333	0.1464	c_3	0.1429	0.2000	0.1184	0.2000	0.1122
Asy.Risk1	1.5769	1.3073		1.2107		Asy.Risk1	1.0238	0.8688		0.7801	
R.R.R.		17.10		23.23		R.R.R.		15.14		23.81	
Asy.Risk2	2.4000		1.2543		1.1919	Asy.Risk2	1.7143		1.0182		0.9455
R.R.R.			47.74		50.34	R.R.R.			40.61		44.84

$n = 9$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 11$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.1111	0.0909	0.0674	0.0732	0.0616	c_1	0.0909	0.0769	0.0592	0.0623	0.0537
c_2	0.1111	0.1111	0.0781	0.1577	0.1162	c_2	0.0909	0.0909	0.0674	0.1260	0.0980
c_3	0.1111	0.1429	0.0950	0.1429	0.0914	c_3	0.0909	0.1111	0.0794	0.1111	0.0771
Asy.Risk1	0.7635	0.6592		0.5793		Asy.Risk1	0.6107	0.5342		0.4622	
R.R.R.		13.66		24.12		R.R.R.		12.52		24.31	
Asy.Risk2	1.3333		0.8585		0.7821	Asy.Risk2	1.0909		0.7430		0.6663
R.R.R.			35.61		41.34	R.R.R.			31.89		38.93

$n = 21$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 51$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.0476	0.0435	0.0371	0.0361	0.0331	c_1	0.0196	0.0189	0.0175	0.0164	0.0158
c_2	0.0476	0.0476	0.0401	0.0613	0.0537	c_2	0.0196	0.0196	0.0182	0.0232	0.0220
c_3	0.0476	0.0526	0.0439	0.0526	0.0435	c_3	0.0196	0.0204	0.0189	0.0204	0.0189
Asy.Risk1	0.3040	0.2755		0.2281		Asy.Risk1	0.1191	0.1117		0.0881	
R.R.R.		9.37		24.97		R.R.R.		6.21		25.99	
Asy.Risk2	0.5714		0.4473		0.3815	Asy.Risk2	0.2353		0.2066		0.1666
R.R.R.			21.71		33.24	R.R.R.			12.20		29.18

Table 3: $p = 4, m = 1$

$n = 5$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 7$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.2000	0.1250	0.0822	0.1250	0.0759	c_1	0.1429	0.1000	0.0690	0.1000	0.0647
c_2	0.2000	0.1667	0.0973	0.1200	0.0927	c_2	0.1429	0.1250	0.0796	0.0883	0.0726
c_3	0.2000	0.2500	0.1222	0.3333	0.2310	c_3	0.1429	0.1667	0.0959	0.2000	0.1559
c_4	0.2000	0.5000	0.1746	1.5000	0.6931	c_4	0.1429	0.2500	0.1259	0.5956	0.3816
Asy.Risk1	3.0752	2.0603		1.5303		Asy.Risk1	1.8508	1.2955		0.9241	
R.R.R.		33.00		50.24		R.R.R.		30.01		50.07	
Asy.Risk2	4.0000		1.8435		1.4655	Asy.Risk2	2.8571		1.4923		1.1116
R.R.R.			53.91		63.36	R.R.R.			47.77		61.10

$n = 9$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 11$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.1111	0.0833	0.0600	0.0833	0.0571	c_1	0.0909	0.0714	0.0533	0.0714	0.0513
c_2	0.1111	0.1000	0.0681	0.0707	0.0602	c_2	0.0909	0.0833	0.0596	0.0593	0.0517
c_3	0.1111	0.1250	0.0798	0.1429	0.1179	c_3	0.0909	0.1000	0.0685	0.1111	0.0949
c_4	0.1111	0.1667	0.0990	0.3497	0.2553	c_4	0.0909	0.1250	0.0819	0.2413	0.1890
Asy.Risk1	1.3436	0.9790		0.6852		Asy.Risk1	1.0585	0.7956		0.5496	
R.R.R.		27.13		49.00		R.R.R.		24.84		48.08	
Asy.Risk2	2.2222		1.2591		0.9083	Asy.Risk2	1.8182		1.0927		0.7730
R.R.R.			43.34		59.13	R.R.R.			39.90		57.49

$n = 21$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 51$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.0476	0.0417	0.0346	0.0417	0.0341	c_1	0.0196	0.0185	0.0170	0.0185	0.0169
c_2	0.0476	0.0455	0.0372	0.0338	0.0311	c_2	0.0196	0.0192	0.0176	0.0154	0.0148
c_3	0.0476	0.0500	0.0404	0.0526	0.0483	c_3	0.0196	0.0200	0.0182	0.0204	0.0197
c_4	0.0476	0.0556	0.0444	0.0879	0.0782	c_4	0.0196	0.0208	0.0189	0.0278	0.0266
Asy.Risk1	0.5127	0.4183		0.2769		Asy.Risk1	0.2016	0.1777		0.1122	
R.R.R.		18.41		45.99		R.R.R.		11.88		44.36	
Asy.Risk2	0.9524		0.6708		0.4526	Asy.Risk2	0.3922		0.3207		0.2055
R.R.R.			29.57		52.47	R.R.R.			18.22		47.59

The risks are analytically calculated except for evaluating $\sum_{i=1}^p E[\log \chi_{n-i+1}^2]$ by Monte Carlo simulation method using 10^5 random numbers. ‘‘R.R.R.’’ under ‘‘Asy.Risk1’’ or ‘‘Asy.Risk2’’ shows the risk reduction rate defined by

$$\text{R.R.R. of } \widehat{\Sigma} = \frac{\text{The risk of } \widehat{\Sigma}^U - \text{The risk of } \widehat{\Sigma}}{\text{The risk of } \widehat{\Sigma}^U} \times 100.$$

It has been observed that $\widehat{\Sigma}^{SDS}$ and $\widehat{\Sigma}^{KG}$ drastically reduce the risk of Σ^U when the population eigenvalues are close to each other. Lin and Perlman (1985) reports that when $\Sigma = \mathbf{I}_p$, R.R.R. of $\widehat{\Sigma}^{SDS}$ often reaches 70%. See also Sugiura and Ishibayashi (1997) for a risk comparison by elaborate simulation. In the situation of the block-wise dispersion, the risk reduction rate of these estimators rarely approaches 50%. Especially when n is as large as 50, the rate is always under 20%. On the other hand, the risk reduction rates of $\widehat{\Sigma}^{MA_1}$ and $\widehat{\Sigma}^{MA_2}$ are constantly over 30% and often reach 50% irrespective of the values of n . It is interesting that $\widehat{\Sigma}^{MA_2}$ always outperforms $\widehat{\Sigma}^{MA_1}$ in view of R.R.R.

3.4 Simulation studies

In this subsection, we evaluate the performance of $\widehat{\Sigma}^{MA_1}$, $\widehat{\Sigma}^{MA_2}$ by Monte Carlo simulation under the situation (31). As we saw in the previous subsection, in view of the asymptotic risks, $\widehat{\Sigma}^{MA_1}$, $\widehat{\Sigma}^{MA_2}$ provide better risk reduction compared to $\widehat{\Sigma}^{SDS}$, $\widehat{\Sigma}^{KG}$. In practical point view, however, it is important to see how largely the population eigenvalues must be dispersed so that

Table 4: $p = 4, m = 2$

$n = 5$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 7$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.2000	0.1250	0.0822	0.1034	0.0762	c_1	0.1429	0.1000	0.0690	0.0820	0.0632
c_2	0.2000	0.1667	0.0973	0.2308	0.1261	c_2	0.1429	0.1250	0.0796	0.1724	0.1055
c_3	0.2000	0.2500	0.1222	0.2000	0.1173	c_3	0.1429	0.1667	0.0959	0.1304	0.0885
c_4	0.2000	0.5000	0.1746	1.0000	0.3988	c_4	0.1429	0.2500	0.1259	0.4286	0.2425
Asy.Risk1	3.0752	2.2687		1.9819		Asy.Risk1	1.8508	1.4334		1.2107	
R.R.R.		26.23		35.55		R.R.R.		22.55		34.59	
Asy.Risk2	4.0000		1.8668		1.7317	Asy.Risk2	2.8571		1.5273		1.3728
R.R.R.			53.33		56.71	R.R.R.			46.54		51.95

$n = 9$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 11$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.1111	0.0833	0.0600	0.0682	0.0546	c_1	0.0909	0.0714	0.0533	0.0586	0.0482
c_2	0.1111	0.1000	0.0681	0.1362	0.0910	c_2	0.0909	0.0833	0.0596	0.1119	0.0798
c_3	0.1111	0.1250	0.0798	0.0980	0.0719	c_3	0.0909	0.1000	0.0685	0.0790	0.0609
c_4	0.1111	0.1667	0.0990	0.2632	0.1727	c_4	0.0909	0.1250	0.0819	0.1872	0.1337
Asy.Risk1	1.3436	1.0774		0.8908		Asy.Risk1	1.0585	0.8700		0.7080	
R.R.R.		19.81		33.70		R.R.R.		17.81		33.11	
Asy.Risk2	2.2222		1.2992		1.1422	Asy.Risk2	1.8182		1.1337		0.9792
R.R.R.			41.54		48.60	R.R.R.			37.65		46.14

$n = 21$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 51$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.0476	0.0417	0.0346	0.0349	0.0330	c_1	0.0196	0.0185	0.0170	0.0162	0.0153
c_2	0.0476	0.0455	0.0372	0.0577	0.0531	c_2	0.0196	0.0192	0.0176	0.0227	0.0211
c_3	0.0476	0.0500	0.0404	0.0410	0.0352	c_3	0.0196	0.0200	0.0182	0.0173	0.0163
c_4	0.0476	0.0556	0.0444	0.0735	0.0615	c_4	0.0196	0.0208	0.0189	0.0248	0.0232
Asy.Risk1	0.5127	0.4477		0.3477		Asy.Risk1	0.2016	0.1857		0.1377	
R.R.R.		12.68		32.18		R.R.R.		7.90		31.73	
Asy.Risk2	0.9524		0.7013		0.5722	Asy.Risk2	0.3922		0.3331		0.2544
R.R.R.			26.36		39.92	R.R.R.			15.06		35.13

Table 5: $p = 4, m = 3$

$n = 4$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 6$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.2500	0.1429	0.0919	0.1071	0.0852	c_1	0.1667	0.1111	0.0749	0.0812	0.0678
c_2	0.2500	0.2000	0.1111	0.2500	0.1670	c_2	0.1667	0.1429	0.0873	0.1667	0.1248
c_3	0.2500	0.3333	0.1449	0.6000	0.2383	c_3	0.1667	0.2000	0.1072	0.3733	0.2028
c_4	0.2500	1.0000	0.2174	1.0000	0.1698	c_4	0.1667	0.3333	0.1461	0.3333	0.1209
Asy.Risk1	4.8592	3.6569		3.4447		Asy.Risk1	2.2985	1.7446		1.5186	
R.R.R.		24.74		29.11		R.R.R.		24.10		33.93	
Asy.Risk2	5.0000		2.0872		1.9697	Asy.Risk2	3.3333		1.6702		1.5097
R.R.R.			58.26		60.61	R.R.R.			49.89		54.71

$n = 8$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 10$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.1250	0.0909	0.0642	0.0660	0.0569	c_1	0.1000	0.0769	0.0565	0.0560	0.0493
c_2	0.1250	0.1111	0.0733	0.1250	0.0999	c_2	0.1000	0.0909	0.0636	0.1000	0.0833
c_3	0.1250	0.1429	0.0870	0.2591	0.1670	c_3	0.1000	0.1111	0.0737	0.1944	0.1385
c_4	0.1250	0.2000	0.1108	0.2000	0.0966	c_4	0.1000	0.1429	0.0896	0.1429	0.0810
Asy.Risk1	1.5538	1.2032		0.9929		Asy.Risk1	1.1828	0.9327		0.7412	
R.R.R.		22.57		36.10		R.R.R.		21.15		37.34	
Asy.Risk2	2.5000		1.3948		1.2111	Asy.Risk2	2.0000		1.1991		1.0067
R.R.R.			44.21		51.56	R.R.R.			40.05		49.66

$n = 20$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$	$n = 50$	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
c_1	0.0500	0.0435	0.0358	0.0326	0.0303	c_1	0.0200	0.0189	0.0172	0.0151	0.0146
c_2	0.0500	0.0476	0.0386	0.0500	0.0455	c_2	0.0200	0.0196	0.0179	0.0200	0.0192
c_3	0.0500	0.0526	0.0421	0.0808	0.0694	c_3	0.0200	0.0204	0.0186	0.0271	0.0256
c_4	0.0500	0.0588	0.0465	0.0588	0.0450	c_4	0.0200	0.0213	0.0193	0.0213	0.0192
Asy.Risk1	0.5385	0.4484		0.3218		Asy.Risk1	0.2069	0.1836		0.1207	
R.R.R.		16.73		40.24		R.R.R.		11.28		41.65	
Asy.Risk2	1.0000		0.7122		0.5395	Asy.Risk2	0.4000		0.3293		0.2236
R.R.R.			28.78		46.05	R.R.R.			17.68		44.11

the use of $\widehat{\Sigma}^{MA_d}$, $d = 1, 2$, is recommended. The convergence speed of the distributions given in Theorem 2, which is an interesting topic by itself, is closely related to this problem.

To see the convergence speed in both distributions and risks, we carried out Monte Carlo Simulation for the two cases $p = 3, m = 1$ and $p = 4, m = 1$. In each case, we took 11 values 1.0, 0.8, 0.6, 0.4, 0.2, 10^{-i} ($i = 1, \dots, 6$) in the convergence parameter β , while α is fixed at 1. We took three different values of n in each case and generated 10^6 random Wishart matrices under given p, n, β . The result is given in Table 6 ($p = 3, m = 1$) and Table 7 ($p = 4, m = 1$). The upper part of each table shows the speed of the distributional convergence in Theorem 2. Note that when $\Xi_1 = \mathbf{I}_m$, $\Xi_2 = \mathbf{I}_{p-m}$, the asymptotic distribution of a diagonal element of $\widehat{\mathbf{W}}_{ss}$, $s = 1, 2$, is a χ^2 distribution. The labels in the tables are given as follows with $\chi_n^2(\alpha)$, $z(\alpha)$ denoting the lower α percentage points of χ^2 distribution with n degrees of freedom and the standard normal distribution, respectively ;

Table 6

$$\begin{aligned} \text{Prob 1a} &= P(\widehat{\mathbf{W}}_{11} \leq \chi_n^2(0.05)), & \text{Prob 1b} &= P(\widehat{\mathbf{W}}_{11} \leq \chi_n^2(0.95)), \\ \text{Prob 2a} &= P((\widehat{\mathbf{W}}_{22})_{11} \leq \chi_{n-1}^2(0.05)), & \text{Prob 2b} &= P((\widehat{\mathbf{W}}_{22})_{11} \leq \chi_{n-1}^2(0.95)), \\ \text{Prob 3a} &= P((\widehat{\mathbf{W}}_{22})_{22} \leq \chi_{n-1}^2(0.05)), & \text{Prob 3b} &= P((\widehat{\mathbf{W}}_{22})_{22} \leq \chi_{n-1}^2(0.95)), \\ \text{Prob 4a} &= P((\widetilde{\mathbf{Z}}_{21})_{11} \leq z(0.05)), & \text{Prob 4b} &= P((\widetilde{\mathbf{Z}}_{21})_{11} \leq z(0.95)), \\ \text{Prob 5a} &= P((\widetilde{\mathbf{Z}}_{21})_{21} \leq z(0.05)), & \text{Prob 5b} &= P((\widetilde{\mathbf{Z}}_{21})_{21} \leq z(0.95)), \end{aligned}$$

Table 7

$$\begin{aligned} \text{Prob 1a} &= P(\widehat{\mathbf{W}}_{11} \leq \chi_n^2(0.05)), & \text{Prob 1b} &= P(\widehat{\mathbf{W}}_{11} \leq \chi_n^2(0.95)), \\ \text{Prob 2a} &= P((\widehat{\mathbf{W}}_{22})_{11} \leq \chi_{n-1}^2(0.05)), & \text{Prob 2b} &= P((\widehat{\mathbf{W}}_{22})_{11} \leq \chi_{n-1}^2(0.95)), \\ \text{Prob 3a} &= P((\widehat{\mathbf{W}}_{22})_{33} \leq \chi_{n-1}^2(0.05)), & \text{Prob 3b} &= P((\widehat{\mathbf{W}}_{22})_{33} \leq \chi_{n-1}^2(0.95)), \\ \text{Prob 4a} &= P((\widetilde{\mathbf{Z}}_{21})_{11} \leq z(0.05)), & \text{Prob 4b} &= P((\widetilde{\mathbf{Z}}_{21})_{11} \leq z(0.95)), \\ \text{Prob 5a} &= P((\widetilde{\mathbf{Z}}_{21})_{31} \leq z(0.05)), & \text{Prob 5b} &= P((\widetilde{\mathbf{Z}}_{21})_{31} \leq z(0.95)). \end{aligned}$$

In the lower part of each table, “Risk 1_*” and “Risk 2_*” show the risks of the corresponding estimator $\widehat{\Sigma}^*$ respectively for L_1 and L_2 . The tables show that

1. The convergence of the diagonal elements of $\widehat{\mathbf{W}}_{ss}$, $s = 1, 2$, is so rapid that when $\beta = 0.1$, the asymptotic distribution already gives a good approximation for the exact distribution. When $\beta = 0.1$, every probability of the diagonal elements is within 0.01 deviation from the exact asymptotic probability.
2. The convergence speed of $\widetilde{\mathbf{Z}}$ is quite slow compared to that of the diagonal elements of $\widehat{\mathbf{W}}_{ss}$, $s = 1, 2$. For a good approximation as above, β must be as small as 10^{-5} or 10^{-6} .
3. The risks also rapidly converge to the asymptotic risks so that $\beta = 0.1$ is small enough to give a good approximation. Actually all the risks in the tables when $\beta = 0.1$ are within the $\pm 5\%$ interval centered at the exact asymptotic risk.
4. The risk of $\widehat{\Sigma}^{MA_d}$, $d = 1, 2$, is always lower than that of the competing estimators. Most notably their superiority in risk is kept even when the population eigenvalues are all equal. It seems that $\widehat{\Sigma}^{MA_d}$, $d = 1, 2$, has robustness to the deviation from the dispersion of the population eigenvalues.

Table 6: $p = 3, m = 1$

$n = 10$	1	0.8	0.6	0.4	0.2	10^{-1}	10^{-2}	10^{-3}	10^{-4}	10^{-5}	10^{-6}	Asymp.
Prob 1a	0.4994	0.3992	0.2814	0.1551	0.0695	0.0534	0.0501	0.0491	0.0507	0.0491	0.0508	0.0500
Prob 2a	0.4091	0.3273	0.2321	0.1321	0.0677	0.0558	0.0516	0.0489	0.0504	0.0495	0.0502	0.0500
Prob 3a	0.4121	0.3302	0.2311	0.1317	0.0684	0.0564	0.0503	0.0499	0.0505	0.0499	0.0518	0.0500
Prob 4a	0.2024	0.1799	0.1502	0.1072	0.0597	0.0385	0.0263	0.0255	0.0294	0.0429	0.0499	0.0500
Prob 5a	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0269	0.0500	0.0500
Prob 1b	0.9700	0.9636	0.9572	0.9531	0.9503	0.9507	0.9499	0.9514	0.9496	0.9497	0.9496	0.9500
Prob 2b	0.9993	0.9985	0.9955	0.9871	0.9695	0.9576	0.9508	0.9498	0.9503	0.9488	0.9498	0.9500
Prob 3b	0.9994	0.9983	0.9957	0.9874	0.9693	0.9582	0.9508	0.9509	0.9496	0.9500	0.9497	0.9500
Prob 4b	0.6174	0.6492	0.6986	0.7671	0.8528	0.8924	0.9236	0.9255	0.9301	0.9451	0.9515	0.9500
Prob 5b	0.4137	0.4673	0.5465	0.6624	0.7937	0.8530	0.8971	0.8994	0.9001	0.9275	0.9504	0.9500
Risk 1_U	0.6769	0.6753	0.6786	0.6779	0.6777	0.6778	0.6784	0.6757	0.6759	0.6758	0.6800	0.6765
Risk 1_SDS	0.4589	0.4611	0.4770	0.5038	0.5409	0.5580	0.5701	0.5687	0.5690	0.5684	0.5727	0.5692
Risk 1_MA1	0.3595	0.3644	0.3824	0.4091	0.4400	0.4553	0.4677	0.4668	0.4677	0.4660	0.4704	0.4676
Risk 2_U	1.1996	1.1997	1.2017	1.1976	1.1983	1.1989	1.1980	1.1966	1.2020	1.1990	1.2021	1.2000
Risk 2_KG	0.7117	0.7132	0.7228	0.7407	0.7641	0.7748	0.7815	0.7812	0.7806	0.7811	0.7839	0.7817
Risk 2_MA2	0.6109	0.6147	0.6255	0.6397	0.6540	0.6625	0.6706	0.6703	0.6704	0.6689	0.6725	0.6591

$n = 20$	1	0.8	0.6	0.4	0.2	10^{-1}	10^{-2}	10^{-3}	10^{-4}	10^{-5}	10^{-6}	Asymp.
Prob 1a	0.6164	0.4550	0.2706	0.1187	0.0574	0.0523	0.0495	0.0506	0.0517	0.0491	0.0498	0.0500
Prob 2a	0.5114	0.3837	0.2285	0.1027	0.0608	0.0537	0.0511	0.0498	0.0505	0.0490	0.0494	0.0500
Prob 3a	0.5081	0.3856	0.2309	0.1043	0.0594	0.0528	0.0511	0.0508	0.0493	0.0513	0.0499	0.0500
Prob 4a	0.2493	0.2196	0.1684	0.1100	0.0560	0.0377	0.0257	0.0264	0.0328	0.0483	0.0513	0.0500
Prob 5a	0.0015	0.0015	0.0008	0.0002	0.0000	0.0000	0.0000	0.0000	0.0003	0.0451	0.0497	0.0500
Prob 1b	0.9767	0.9661	0.9595	0.9543	0.9515	0.9484	0.9513	0.9500	0.9498	0.9498	0.9494	0.9500
Prob 2b	0.9995	0.9984	0.9936	0.9816	0.9631	0.9547	0.9513	0.9499	0.9498	0.9511	0.9500	0.9500
Prob 3b	0.9994	0.9983	0.9940	0.9815	0.9623	0.9552	0.9520	0.9516	0.9499	0.9501	0.9505	0.9500
Prob 4b	0.5542	0.6008	0.6689	0.7653	0.8592	0.8975	0.9200	0.9257	0.9334	0.9499	0.9508	0.9500
Prob 5b	0.3123	0.3793	0.4993	0.6566	0.8026	0.8574	0.8953	0.8998	0.8987	0.9457	0.9503	0.9500
Risk 1_U	0.3178	0.3179	0.3181	0.3171	0.3183	0.3178	0.3177	0.3175	0.3177	0.3171	0.3176	0.3164
Risk 1_SDS	0.2363	0.2390	0.2486	0.2628	0.2767	0.2802	0.2829	0.2830	0.2833	0.2827	0.2832	0.2819
Risk 1_MA1	0.1880	0.1923	0.2023	0.2115	0.2200	0.2236	0.2261	0.2262	0.2267	0.2260	0.2262	0.2251
Risk 2_U	0.5995	0.6011	0.6008	0.5992	0.5999	0.6006	0.5992	0.5987	0.6005	0.6003	0.6011	0.6000
Risk 2_KG	0.4085	0.4117	0.4226	0.4384	0.4530	0.4568	0.4595	0.4598	0.4600	0.4593	0.4594	0.4598
Risk 2_MA2	0.3563	0.3606	0.3674	0.3706	0.3744	0.3775	0.3792	0.3794	0.3801	0.3797	0.3793	0.3745

$n = 50$	1	0.8	0.6	0.4	0.2	10^{-1}	10^{-2}	10^{-3}	10^{-4}	10^{-5}	10^{-6}	Asymp.
Prob 1a	0.7358	0.4769	0.2076	0.0793	0.0532	0.0503	0.0484	0.0506	0.0506	0.0485	0.0501	0.0500
Prob 2a	0.6110	0.4089	0.1725	0.0720	0.0549	0.0511	0.0512	0.0498	0.0505	0.0489	0.0499	0.0500
Prob 3a	0.6079	0.4100	0.1737	0.0732	0.0564	0.0513	0.0495	0.0484	0.0487	0.0489	0.0504	0.0500
Prob 4a	0.2992	0.2529	0.1788	0.1042	0.0541	0.0368	0.0271	0.0274	0.0411	0.0493	0.0500	0.0500
Prob 5a	0.0200	0.0109	0.0031	0.0002	0.0000	0.0000	0.0000	0.0000	0.0072	0.0490	0.0506	0.0500
Prob 1b	0.9823	0.9707	0.9606	0.9532	0.9511	0.9499	0.9485	0.9503	0.9489	0.9500	0.9505	0.9500
Prob 2b	0.9995	0.9979	0.9883	0.9696	0.9567	0.9519	0.9503	0.9511	0.9498	0.9506	0.9498	0.9500
Prob 3b	0.9996	0.9977	0.9889	0.9705	0.9557	0.9528	0.9495	0.9499	0.9500	0.9506	0.9498	0.9500
Prob 4b	0.5050	0.5599	0.6582	0.7701	0.8610	0.8967	0.9231	0.9266	0.9408	0.9502	0.9495	0.9500
Prob 5b	0.2273	0.3156	0.4805	0.6648	0.8100	0.8613	0.8960	0.8993	0.9066	0.9494	0.9498	0.9500
Risk 1_U	0.1223	0.1226	0.1227	0.1229	0.1228	0.1228	0.1226	0.1223	0.1228	0.1221	0.1230	0.1236
Risk 1_SDS	0.1006	0.1026	0.1074	0.1117	0.1137	0.1143	0.1145	0.1143	0.1148	0.1140	0.1149	0.1155
Risk 1_MA1	0.0814	0.0843	0.0867	0.0871	0.0882	0.0888	0.0891	0.0891	0.0896	0.0887	0.0895	0.0901
Risk 2_U	0.2391	0.2399	0.2400	0.2401	0.2402	0.2404	0.2403	0.2391	0.2405	0.2389	0.2406	0.2400
Risk 2_KG	0.1874	0.1906	0.1982	0.2049	0.2079	0.2087	0.2091	0.2086	0.2096	0.2083	0.2097	0.2093
Risk 2_MA2	0.1641	0.1673	0.1669	0.1643	0.1649	0.1654	0.1658	0.1656	0.1665	0.1650	0.1663	0.1647

Because of the robustness, $\hat{\Sigma}^{MA_d}$, $d = 1, 2$, seem to be useful for various applications. Now as the last topic in this section, apart from a decision-theoretic approach, we evaluate these new estimators' performance in discriminant analysis. We use a well-known example of Fisher's iris data. The data consists of 50 samples from each of the three groups(species) with 4-dimensional variable (x_1 :sepal length(cm), x_2 :sepal width(cm), x_3 :petal length(cm), x_4 :petal width(cm)). We

Table 7: $p = 4, m = 1$

$n = 11$	1	0.8	0.6	0.4	0.2	10^{-1}	10^{-2}	10^{-3}	10^{-4}	10^{-5}	10^{-6}	Asymp.
Prob 1a	0.5852	0.4760	0.3396	0.1881	0.0751	0.0554	0.0517	0.0497	0.0495	0.0498	0.0489	0.0500
Prob 2a	0.3156	0.2620	0.1937	0.1166	0.0660	0.0557	0.0505	0.0494	0.0510	0.0501	0.0492	0.0500
Prob 3a	0.3146	0.2620	0.1920	0.1172	0.0651	0.0560	0.0515	0.0503	0.0496	0.0494	0.0503	0.0500
Prob 4a	0.1955	0.1794	0.1533	0.1124	0.0650	0.0431	0.0283	0.0284	0.0315	0.0449	0.0507	0.0500
Prob 5a	0.1312	0.1206	0.1005	0.0685	0.0342	0.0205	0.0131	0.0131	0.0172	0.0391	0.0497	0.0500
Prob 1b	0.9761	0.9676	0.9610	0.9547	0.9521	0.9510	0.9500	0.9510	0.9495	0.9508	0.9501	0.9500
Prob 2b	0.9986	0.9977	0.9948	0.9863	0.9690	0.9581	0.9505	0.9510	0.9507	0.9509	0.9493	0.9500
Prob 3b	0.9985	0.9979	0.9947	0.9861	0.9681	0.9570	0.9509	0.9515	0.9499	0.9508	0.9508	0.9500
Prob 4b	0.6525	0.6774	0.7152	0.7789	0.8576	0.8979	0.9251	0.9276	0.9310	0.9454	0.9501	0.9500
Prob 5b	0.5899	0.6184	0.6607	0.7327	0.8304	0.8751	0.9091	0.9115	0.9185	0.9399	0.9508	0.9500
Risk 1_U	1.0566	1.0583	1.0552	1.0592	1.0577	1.0583	1.0603	1.0544	1.0574	1.0573	1.0559	1.0585
Risk 1_SDS	0.6514	0.6572	0.6714	0.7092	0.7558	0.7781	0.7954	0.7920	0.7943	0.7942	0.7927	0.7956
Risk 1_MA1	0.4064	0.4154	0.4367	0.4738	0.5104	0.5295	0.5485	0.5471	0.5484	0.5478	0.5468	0.5496
Risk 2_U	1.8199	1.8213	1.8147	1.8170	1.8175	1.8199	1.8210	1.8147	1.8206	1.8180	1.8176	1.8182
Risk 2_KG	1.0173	1.0214	1.0291	1.0493	1.0749	1.0876	1.0939	1.0921	1.0929	1.0926	1.0915	1.0927
Risk 2_MA2	0.5967	0.6075	0.6326	0.6767	0.7268	0.7516	0.7728	0.7724	0.7737	0.7733	0.7719	0.7730

$n = 21$	1	0.8	0.6	0.4	0.2	10^{-1}	10^{-2}	10^{-3}	10^{-4}	10^{-5}	10^{-6}	Asymp.
Prob 1a	0.7030	0.5419	0.3317	0.1428	0.0601	0.0532	0.0503	0.0505	0.0498	0.0495	0.0509	0.0500
Prob 2a	0.4043	0.3183	0.2017	0.0985	0.0579	0.0547	0.0508	0.0495	0.0492	0.0505	0.0505	0.0500
Prob 3a	0.3995	0.3222	0.2019	0.0975	0.0581	0.0527	0.0507	0.0504	0.0496	0.0497	0.0493	0.0500
Prob 4a	0.2413	0.2156	0.1748	0.1172	0.0601	0.0421	0.0297	0.0292	0.0344	0.0480	0.0503	0.0500
Prob 5a	0.1720	0.1533	0.1185	0.0711	0.0331	0.0201	0.0141	0.0137	0.0211	0.0484	0.0505	0.0500
Prob 1b	0.9830	0.9737	0.9627	0.9557	0.9502	0.9506	0.9503	0.9504	0.9497	0.9505	0.9505	0.9500
Prob 2b	0.9989	0.9977	0.9929	0.9809	0.9617	0.9548	0.9507	0.9501	0.9488	0.9500	0.9514	0.9500
Prob 3b	0.9988	0.9975	0.9935	0.9805	0.9628	0.9549	0.9509	0.9502	0.9501	0.9496	0.9487	0.9500
Prob 4b	0.5985	0.6278	0.6881	0.7757	0.8657	0.8998	0.9262	0.9272	0.9339	0.9476	0.9494	0.9500
Prob 5b	0.5303	0.5632	0.6291	0.7282	0.8368	0.8794	0.9118	0.9130	0.9219	0.9479	0.9504	0.9500
Risk 1_U	0.5121	0.5136	0.5135	0.5116	0.5115	0.5128	0.5110	0.5127	0.5115	0.5109	0.5119	0.5127
Risk 1_SDS	0.3503	0.3552	0.3677	0.3871	0.4056	0.4134	0.4167	0.4183	0.4172	0.4169	0.4177	0.4183
Risk 1_MA1	0.2241	0.2315	0.2461	0.2568	0.2650	0.2715	0.2759	0.2772	0.2759	0.2764	0.2765	0.2769
Risk 2_U	0.9512	0.9514	0.9537	0.9503	0.9516	0.9521	0.9477	0.9535	0.9520	0.9501	0.9505	0.9524
Risk 2_KG	0.6059	0.6109	0.6233	0.6429	0.6607	0.6669	0.6692	0.6708	0.6700	0.6695	0.6707	0.6708
Risk 2_MA2	0.3510	0.3622	0.3861	0.4114	0.4326	0.4433	0.4516	0.4532	0.4521	0.4524	0.4525	0.4526

$n = 51$	1	0.8	0.6	0.4	0.2	10^{-1}	10^{-2}	10^{-3}	10^{-4}	10^{-5}	10^{-6}	Asymp.
Prob 1a	0.8209	0.5805	0.2691	0.0916	0.0533	0.0492	0.0498	0.0504	0.0501	0.0504	0.0500	0.0500
Prob 2a	0.5101	0.3626	0.1647	0.0721	0.0560	0.0522	0.0501	0.0502	0.0501	0.0504	0.0500	0.0500
Prob 3a	0.5098	0.3610	0.1669	0.0722	0.0555	0.0533	0.0500	0.0507	0.0479	0.0498	0.0506	0.0500
Prob 4a	0.2912	0.2595	0.1878	0.1118	0.0604	0.0415	0.0303	0.0291	0.0403	0.0507	0.0491	0.0500
Prob 5a	0.2191	0.1863	0.1307	0.0689	0.0308	0.0196	0.0133	0.0148	0.0313	0.0501	0.0499	0.0500
Prob 1b	0.9891	0.9762	0.9649	0.9548	0.9507	0.9501	0.9504	0.9501	0.9504	0.9505	0.9497	0.9500
Prob 2b	0.9992	0.9970	0.9889	0.9700	0.9573	0.9526	0.9502	0.9498	0.9503	0.9507	0.9504	0.9500
Prob 3b	0.9990	0.9973	0.9891	0.9712	0.9565	0.9521	0.9499	0.9513	0.9503	0.9506	0.9492	0.9500
Prob 4b	0.5383	0.5836	0.6703	0.7803	0.8683	0.9022	0.9272	0.9286	0.9411	0.9494	0.9503	0.9500
Prob 5b	0.4666	0.5129	0.6101	0.7334	0.8386	0.8789	0.9081	0.9153	0.9312	0.9496	0.9503	0.9500
Risk 1_U	0.2018	0.2022	0.2019	0.2017	0.2019	0.2017	0.2020	0.2017	0.2020	0.2023	0.2018	0.2016
Risk 1_SDS	0.1566	0.1592	0.1658	0.1721	0.1758	0.1768	0.1780	0.1777	0.1780	0.1783	0.1779	0.1777
Risk 1_MA1	0.1037	0.1083	0.1109	0.1088	0.1104	0.1113	0.1125	0.1124	0.1123	0.1124	0.1125	0.1122
Risk 2_U	0.3923	0.3939	0.3920	0.3916	0.3920	0.3924	0.3927	0.3919	0.3931	0.3929	0.3920	0.3922
Risk 2_KG	0.2896	0.2938	0.3038	0.3127	0.3179	0.3194	0.3208	0.3203	0.3211	0.3215	0.3208	0.3207
Risk 2_MA2	0.1785	0.1867	0.1943	0.1959	0.2010	0.2033	0.2057	0.2055	0.2054	0.2056	0.2059	0.2055

downloaded the data from the website <http://www-unix.oit.umass.edu/~statdata>. We let $\mathbf{x}_j^{(i)}$, $i = 1, 2, 3$, $j = 1, \dots, 50$ denote the j th sample in the i th group. The estimator to be tested are the traditional estimators $\hat{\Sigma}^U$, $\hat{\Sigma}^{SDS}$, $\hat{\Sigma}^{KG}$ and the new estimators $\hat{\Sigma}^{MA_1}$, $\hat{\Sigma}^{MA_2}$ which are formulated under the condition $p = 4, m = 1$.

We carry out cross validations. Suppose a learning data set $\mathbf{y}_j^{(i)}$, $j = 1, \dots, N$, is chosen from

Table 8: 10-sample-set

Learning Data Set	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
1	82.50	83.33	83.33	81.67	82.50
2	85.83	85.00	85.00	85.00	85.00
3	82.50	82.50	82.50	82.50	82.50
4	81.67	83.33	82.50	85.83	84.17
5	76.67	77.50	77.50	79.17	79.17
Average	81.83	82.33	82.17	82.83	82.67

Table 9: 5-sample-set

Learning Data Set	$\widehat{\Sigma}^U$	$\widehat{\Sigma}^{SDS}$	$\widehat{\Sigma}^{KG}$	$\widehat{\Sigma}^{MA_1}$	$\widehat{\Sigma}^{MA_2}$
1	66.67	71.85	68.89	75.56	75.56
2	78.52	80.00	78.52	85.19	82.96
3	41.48	41.48	41.48	44.44	42.96
4	43.70	46.67	45.93	53.33	50.37
5	88.89	88.15	88.89	92.59	90.37
6	73.33	78.52	77.78	89.63	88.15
7	64.44	68.89	67.41	73.33	71.85
8	73.33	75.56	72.59	82.96	79.26
9	73.33	75.56	72.59	82.96	79.26
10	69.63	72.59	71.85	82.22	77.78
Average	67.33	69.93	68.59	76.22	73.85

the i th group, $i = 1, 2, 3$. Estimates for the population covariance matrix of the i th group are calculated from $\widehat{\Sigma}^U$, $\widehat{\Sigma}^{SDS}$, $\widehat{\Sigma}^{KG}$, $\widehat{\Sigma}^{MA_1}$, $\widehat{\Sigma}^{MA_2}$ based on

$$\mathbf{A}^{(i)} = \sum_{j=1}^N (\mathbf{y}_j^{(i)} - \bar{\mathbf{y}}^{(i)})(\mathbf{y}_j^{(i)} - \bar{\mathbf{y}}^{(i)})',$$

where $\bar{\mathbf{y}}^{(i)} = N^{-1} \sum_{j=1}^N \mathbf{y}_j^{(i)}$. As a discriminant function, we use a Mahalanobis distance based on each estimates $\widehat{\Sigma}^U(\mathbf{A}^{(i)})$, $\widehat{\Sigma}^{SDS}(\mathbf{A}^{(i)})$, $\widehat{\Sigma}^{KG}(\mathbf{A}^{(i)})$, $\widehat{\Sigma}^{MA_1}(\mathbf{A}^{(i)})$, $\widehat{\Sigma}^{MA_2}(\mathbf{A}^{(i)})$, that is, for a test data \mathbf{x}

$$MD_i^* = (\mathbf{x} - \bar{\mathbf{y}}^{(i)})' \widehat{\Sigma}^*(\mathbf{A}^{(i)})^{-1} (\mathbf{x} - \bar{\mathbf{y}}^{(i)}), \quad i = 1, 2, 3.$$

The eigenvalues of the covariance matrix within each group is as follows;

$$\begin{aligned} \text{Group 1: } & (0.234, 0.039, 0.027, 0.009), \\ \text{Group 2: } & (0.482, 0.075, 0.056, 0.011), \\ \text{Group 3: } & (0.688, 0.107, 0.057, 0.036). \end{aligned} \tag{34}$$

We observe that 1) in each group, the largest eigenvalue are about 6 times as large as the second largest eigenvalue, 2) the second largest eigenvalue is about 3–7 times as large as the smallest eigenvalue. We are interested in the performance of Σ^{MA_d} , $d = 1, 2$, with the population

eigenvalues in (34) which are considered as a deviation from (∞, c, c, c) , the ideal eigenvalues for Σ^{MA_i} , $i = 1, 2$.

We made three types of cross validations.

1. Leave-one-out: For a chosen (i, j) , $i = 1, 2, 3$, $j = 1, \dots, 50$, leave $\mathbf{x}_j^{(i)}$ out from the whole data to be a test data, and use the rest as a learning data set. We repeat this trial for every possible (i, j) . Consequently 150 trials were carried out.
2. 10-sample-set: First choose $\mathbf{x}_1^{(i)}, \dots, \mathbf{x}_{10}^{(i)}$, $i = 1, 2, 3$, as a learning data set and use all the rest as a test data. Next use $\mathbf{x}_{11}^{(i)}, \dots, \mathbf{x}_{20}^{(i)}$, $i = 1, 2, 3$, as a learning data set and the others as a test data. Repeatedly change a learning data set until every data is used once as a learning data. Totally we carried out $600 (= 120 \times 5)$ trials.
3. 5-sample-set: First choose $\mathbf{x}_1^{(i)}, \dots, \mathbf{x}_5^{(i)}$, $i = 1, 2, 3$, as a learning data set and use all the rest as a test data. Next use $\mathbf{x}_6^{(i)}, \dots, \mathbf{x}_{10}^{(i)}$, $i = 1, 2, 3$, as a learning data set and the others as a test data. Repeatedly change a learning data set until every data is used once as a learning data. Totally we carried out $1350 (= 135 \times 10)$ trials.

We summarize the result on the correct classification percentage (“C.C.P.” for abbreviation) of each discriminant function.

1. Leave-one-out: All the discriminant functions returned the same classification for every test data and scored 96.67% of C.C.P. The misclassification occurred at the sample $\mathbf{x}_{19}^{(2)}$, $\mathbf{x}_{21}^{(2)}$, $\mathbf{x}_{23}^{(2)}$, $\mathbf{x}_{34}^{(2)}$, $\mathbf{x}_{32}^{(3)}$. With as much as 49 learning data, all the discriminant functions work quite correctly and make no differences among the functions.
2. 10-sample-set: See Table 8 for the C.C.P. in each learning data set and the average. Depending on the learning data set, different discriminant functions records the best C.C.P, but the margins are small and negligible. It seems that even 10-sample-learning set is too large to differentiate the functions.
3. 5-sample-set: See Table 9 for the C.C.P. in each learning data set and the average. In every learning data set, the functions based on $\widehat{\Sigma}^{MA_d}$, $d = 1, 2$, outperform the other functions. Especially $\widehat{\Sigma}^{MA_1}$ always keeps the highest C.C.P. In total, $\widehat{\Sigma}^{MA_1}$ and $\widehat{\Sigma}^{MA_2}$ record better C.C.P. than $\widehat{\Sigma}^U$ by 8.89% and 6.52% respectively, while the margins of $\widehat{\Sigma}^{SDS}$ and $\widehat{\Sigma}^{KG}$ over $\widehat{\Sigma}^U$ are respectively 2.60% and 1.26%.

A Appendix

A.1 Proof of Lemma 1

In the following, c_i ($i = 1, \dots, 7$) represents some constant independent of α, β .

The random variables $\mathbf{l} = (l_1, \dots, l_p)$ and $\widetilde{\mathbf{G}} = \mathbf{\Gamma}'\mathbf{G}$ have the following joint density function with respect to the product measure between Lebesgue measure on \mathcal{L} and the invariant probability μ on $\mathcal{O}^+(p)$.

$$c_1 \prod_{i=1}^p \lambda_i^{-\frac{n}{2}} \prod_{i=1}^p l_i^{\frac{n-p-1}{2}} \prod_{j < i} (l_j - l_i) \operatorname{etr} \left(-\frac{1}{2} \widetilde{\mathbf{G}} \mathbf{L} \widetilde{\mathbf{G}}' \mathbf{\Lambda}^{-1} \right).$$

We have

$$\begin{aligned}
E[x(\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta)] &= E[x(\boldsymbol{\Gamma}\widetilde{\mathbf{G}}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta)] \\
&= c_1 \prod_{i=1}^p \lambda_i^{-\frac{n}{2}} \int_{\mathcal{L}} \int_{\mathcal{O}(p)^+} x(\boldsymbol{\Gamma}\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) \prod_{i=1}^p l_i^{\frac{n-p-1}{2}} \prod_{j<i} (l_j - l_i) \\
&\quad \times \text{etr} \left(-\frac{1}{2} \mathbf{G}\mathbf{L}\mathbf{G}'\boldsymbol{\Lambda}^{-1} \right) d\mu(\mathbf{G})d\mathbf{l}.
\end{aligned}$$

Using the finite open cover $\mathcal{O}^{(\tau)}$, $\tau = 0, \dots, T$, in Subsection 2.1, we have

$$E[x(\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta)] = \sum_{\tau=0}^T I_{\tau}, \quad (35)$$

where

$$\begin{aligned}
I_{\tau} &= c_1 \prod_{i=1}^p \lambda_i^{-\frac{n}{2}} \int_{\mathcal{L}} \int_{\mathcal{O}^+(p)} \iota_{\tau}(\mathbf{G}) x(\boldsymbol{\Gamma}\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) \prod_{i=1}^p l_i^{\frac{n-p-1}{2}} \prod_{j<i} (l_j - l_i) \\
&\quad \times \text{etr} \left(-\frac{1}{2} \mathbf{G}\mathbf{L}\mathbf{G}'\boldsymbol{\Lambda}^{-1} \right) d\mu(\mathbf{G})d\mathbf{l},
\end{aligned}$$

First we consider I_0 . Let M denote the support of ι_0 . From (19),

$$\begin{aligned}
|I_0| &\leq c_1 \prod_{i=1}^p \lambda_i^{-\frac{n}{2}} \int_{\mathcal{L}} \int_M |x(\boldsymbol{\Gamma}\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta)| \prod_{i=1}^p l_i^{\frac{n-p-1}{2}} \prod_{j<i} (l_j - l_i) \\
&\quad \times \text{etr} \left(-\frac{1}{2} \mathbf{G}\mathbf{L}\mathbf{G}'\boldsymbol{\Lambda}^{-1} \right) d\mu(\mathbf{G})d\mathbf{l} \\
&\leq c_1 b \prod_{i=1}^p \lambda_i^{-\frac{n}{2}} \int_{\mathcal{L}} \int_M \prod_{i=1}^p l_i^{\frac{n-p-1}{2}} \prod_{j<i} (l_j - l_i) \text{etr} \left(-\frac{1}{2} \mathbf{G}\mathbf{L}\mathbf{G}'\bar{\boldsymbol{\Lambda}}^{-1} \right) d\mu(\mathbf{G})d\mathbf{l} \\
&= c_2 P \left(\widetilde{\mathbf{G}} \in M \mid \boldsymbol{\Sigma} = \boldsymbol{\Gamma}\bar{\boldsymbol{\Lambda}}\boldsymbol{\Gamma}' \right), \quad (36)
\end{aligned}$$

where $\bar{\boldsymbol{\Lambda}} = (1 - 2a)^{-1}\boldsymbol{\Lambda}$. Note $\mathcal{O}^+(p) \setminus M$ is an open set including $\mathcal{O}(m, p - m)$, hence by 2 of Theorem 1, $\lim_{\beta/\alpha \rightarrow 0} P \left(\widetilde{\mathbf{G}} \in \mathcal{O}^+(p) \setminus M \mid \boldsymbol{\Sigma} = \boldsymbol{\Gamma}\bar{\boldsymbol{\Lambda}}\boldsymbol{\Gamma}' \right) = 1$, which means

$$P \left(\widetilde{\mathbf{G}} \in M \mid \boldsymbol{\Sigma} = \boldsymbol{\Gamma}\bar{\boldsymbol{\Lambda}}\boldsymbol{\Gamma}' \right) \rightarrow 0$$

as $\beta/\alpha \rightarrow 0$. Therefore

$$\lim_{\beta/\alpha \rightarrow 0} I_0 = 0. \quad (37)$$

Now we focus ourselves on I_{τ} , $\tau = 1, \dots, T$. Since μ is invariant and the support of $\iota_{\tau}(\mathbf{G})$ is contained in $\mathcal{O}^{(\tau)}$, we have

$$\begin{aligned}
I_{\tau} &= c_1 \prod_{i=1}^p \lambda_i^{-\frac{n}{2}} \int_{\mathcal{L}} \int_V \iota_{\tau}(\mathbf{H}^{(\tau)}\mathbf{G}) x(\boldsymbol{\Gamma}\mathbf{H}^{(\tau)}\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) \prod_{i=1}^p l_i^{\frac{n-p-1}{2}} \prod_{j<i} (l_j - l_i) \\
&\quad \times \text{etr} \left(-\frac{1}{2} \mathbf{H}^{(\tau)}\mathbf{G}\mathbf{L}\mathbf{G}'\mathbf{H}^{(\tau)'}\boldsymbol{\Lambda}^{-1} \right) d\mu(\mathbf{G})d\mathbf{l}.
\end{aligned}$$

We want to express the integral with respect to $d\mu(\mathbf{G})$ in terms of the local coordinates \mathbf{u} on U . It is well known that the invariant measure $d\mu(\mathbf{G})$ has the exterior differential form expression

$$c_3 \bigwedge_{i>j} \mathbf{g}'_j d\mathbf{g}_i, \quad (38)$$

where \mathbf{g}_i is the i th column of \mathbf{G} . Substituting the differential

$$\begin{aligned} dg_{ij} &= du_{ij}, \quad i > j, \\ dg_{ij} &= \sum_{k>l} \frac{\partial g_{ij}}{\partial u_{kl}} du_{kl}, \quad i \leq j, \end{aligned}$$

into (38) and taking the wedge product of the terms, we see that

$$\bigwedge_{i>j} \mathbf{g}'_j d\mathbf{g}_i = \pm J^*(\mathbf{u}) \bigwedge_{i>j} du_{ij},$$

where $J^*(\mathbf{u})$ is the Jacobian expressing the Radon-Nikodym derivative of the measure on U induced from the invariant measure on $\mathcal{O}^+(p)$ with respect to the Lebesgue measure on $R^{\frac{p(p-1)}{2}}$. An explicit form of $J^*(\mathbf{u})$ for small dimension p is discussed in Appendix B in Takemura and Sheena (2005). Since $J^*(\mathbf{u})$ is a C^∞ function on \bar{U} , it is bounded and has a finite limit as $\mathbf{u} \rightarrow \mathbf{0}$. By the change of variables $(\mathbf{l}, \mathbf{G}) \rightarrow (\mathbf{l}, \mathbf{u})$, I_τ is written as

$$\begin{aligned} I_\tau &= c_4 \prod_{i=1}^p \lambda_i^{-\frac{n}{2}} \int_{\mathcal{L}} \int_U \iota_\tau(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u})) x(\Gamma \mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}), \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) \prod_{i=1}^p l_i^{\frac{n-p-1}{2}} \prod_{j<i} (l_j - l_i) \\ &\quad \times \text{etr} \left(-\frac{1}{2} \mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}) \mathbf{L} \mathbf{G}'(\mathbf{u}) \mathbf{H}^{(\tau)'} \boldsymbol{\Lambda}^{-1} \right) J^*(\mathbf{u}) d\mathbf{u} d\mathbf{l}, \end{aligned}$$

Consider further coordinate transformation $(\mathbf{l}, \mathbf{u}) \rightarrow (\mathbf{d}, \mathbf{q})$ for each τ . Notice

$$\prod_{i=1}^p l_i^{\frac{n-p-1}{2}} = \left(\prod_{i=1}^p d_j^{\frac{n-p-1}{2}} \right) \alpha^{\frac{m(n-p-1)}{2}} \beta^{\frac{(p-m)(n-p-1)}{2}}, \quad (39)$$

$$\begin{aligned} \prod_{j<i} (l_j - l_i) &= \alpha^{\frac{m(m-1)}{2}} \beta^{\frac{(p-m)(p-m-1)}{2}} \prod_{j \leq m < i} (\alpha d_j - \beta d_i) \prod_{j < i \leq m} (d_j - d_i) \prod_{m < j < i} (d_j - d_i) \\ &= \prod_{j \leq m < i} \left(1 - \frac{\beta d_i}{\alpha d_j} \right) \prod_{j < i \leq m} (d_j - d_i) \prod_{m < j < i} (d_j - d_i) \prod_{j=1}^m d_j^{p-m} \\ &\quad \times \alpha^{m(p-m) + \frac{m(m-1)}{2}} \beta^{\frac{(p-m)(p-m-1)}{2}}, \end{aligned} \quad (40)$$

and

$$\begin{aligned} &\text{tr} \mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}) \mathbf{L} \mathbf{G}'(\mathbf{u}) \mathbf{H}^{(\tau)'} \boldsymbol{\Lambda}^{-1} \\ &= \text{tr} \left\{ \begin{pmatrix} \mathbf{H}_1^{(\tau)} \mathbf{G}_{11}(\mathbf{u}) & \mathbf{H}_1^{(\tau)} \mathbf{G}_{12}(\mathbf{u}) \\ \mathbf{H}_2^{(\tau)} \mathbf{G}_{21}(\mathbf{u}) & \mathbf{H}_2^{(\tau)} \mathbf{G}_{22}(\mathbf{u}) \end{pmatrix} \text{diag}(l_1, \dots, l_p) \right. \\ &\quad \left. \times \begin{pmatrix} \mathbf{G}'_{11}(\mathbf{u}) \mathbf{H}_1^{(\tau)'} & \mathbf{G}'_{21}(\mathbf{u}) \mathbf{H}_2^{(\tau)'} \\ \mathbf{G}'_{12}(\mathbf{u}) \mathbf{H}_1^{(\tau)'} & \mathbf{G}'_{22}(\mathbf{u}) \mathbf{H}_2^{(\tau)'} \end{pmatrix} \text{diag}(\lambda_1^{-1}, \dots, \lambda_p^{-1}) \right\} \\ &= \text{tr} \mathbf{H}_1^{(\tau)} \mathbf{G}_{11}(\mathbf{u}) \mathbf{D}_1 \mathbf{G}'_{11}(\mathbf{u}) \mathbf{H}_1^{(\tau)'} \boldsymbol{\Xi}_1^{-1} + \text{tr} \mathbf{H}_2^{(\tau)} \mathbf{G}_{22}(\mathbf{u}) \mathbf{D}_2 \mathbf{G}'_{22}(\mathbf{u}) \mathbf{H}_2^{(\tau)'} \boldsymbol{\Xi}_2^{-1} \\ &\quad + \text{tr} \mathbf{Q}_{21} \mathbf{Q}'_{21} + \alpha^{-1} \beta \text{tr} \mathbf{H}_1^{(\tau)} \mathbf{G}_{12}(\mathbf{u}) \mathbf{D}_2 \mathbf{G}'_{12}(\mathbf{u}) \mathbf{H}_1^{(\tau)'} \boldsymbol{\Xi}_1^{-1}, \end{aligned} \quad (41)$$

where \mathbf{u} is actually the abbreviation for $\mathbf{u}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)$ defined by (10). For notational simplicity we use the same abbreviation $\mathbf{u} = \mathbf{u}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)$ for the rest of this proof. From (12), (16), (39), (40) and (41), we have

$$I_\tau = c_5 \int_{R^{p(p-1)/2}} \int_{R_+^p} \iota_\tau(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u})) x(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \boldsymbol{\Gamma}, \mathbf{H}^{(\tau)}) h(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta) d\mathbf{d}d\mathbf{q}$$

where $R_+^p = \{\mathbf{d} \mid d_i > 0, i = 1, \dots, p\}$ and $h(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)$ is defined as follows;

$$\begin{aligned} h(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta) &= I(\mathbf{u} \in U) J^*(\mathbf{u}) I(\mathbf{d}_1 \in \mathcal{D}_1, \mathbf{d}_2 \in \mathcal{D}_2, (\mathbf{d}_1, \mathbf{d}_2) \in \mathcal{D}_3) \\ &\times \prod_{i=1}^m d_i^{\frac{n-m-1}{2}} \prod_{i=m+1}^p d_i^{\frac{n-p-1}{2}} \prod_{j < i \leq m} (d_j - d_i) \prod_{m < j < i} (d_j - d_i) \prod_{j \leq m < i} \left(1 - \frac{\beta d_i}{\alpha d_j}\right) \\ &\times \exp\left(-\frac{1}{2} \operatorname{tr} \sum_{s=1}^2 \mathbf{H}_s^{(\tau)} \mathbf{G}_{ss}(\mathbf{u}) \mathbf{D}_s \mathbf{G}'_{ss}(\mathbf{u}) \mathbf{H}_s^{(\tau)'} \boldsymbol{\Xi}_s^{-1}\right) \\ &\times \operatorname{etr}\left(-\frac{1}{2} \mathbf{Q}_{21} \mathbf{Q}'_{21}\right) \times \operatorname{etr}\left(-\frac{\beta}{2\alpha} \mathbf{H}_1^{(\tau)} \mathbf{G}_{12}(\mathbf{u}) \mathbf{D}_2 \mathbf{G}'_{12}(\mathbf{u}) \mathbf{H}_1^{(\tau)'} \boldsymbol{\Xi}_1^{-1}\right). \end{aligned}$$

We will show that

$$\iota_\tau(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u})) x(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \boldsymbol{\Gamma}, \mathbf{H}^{(\tau)}) h(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)$$

is bounded in (α, β) . First $I(\mathbf{u} \in U) J^*(\mathbf{u}) \leq K$ for some $K (> 0)$ since $J^*(\mathbf{u})$ is bounded on the compact set \bar{U} . Clearly

$$0 \leq I(\mathbf{d}_1 \in \mathcal{D}_1, \mathbf{d}_2 \in \mathcal{D}_2, (\mathbf{d}_1, \mathbf{d}_2) \in \mathcal{D}_3) \prod_{j \leq m < i} \left(1 - \frac{\beta d_i}{\alpha d_j}\right) \leq 1.$$

From the condition (19), we have

$$\begin{aligned} |x(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \boldsymbol{\Gamma}, \mathbf{H}^{(\tau)})| &= |x(\boldsymbol{\Gamma} \mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}), \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta)| \\ &\leq b \operatorname{etr}(a \mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}) \mathbf{L} \mathbf{G}'(\mathbf{u}) \mathbf{H}^{(\tau)'} \boldsymbol{\Lambda}^{-1}) \text{ a.e. in } (\mathbf{d}, \mathbf{q}). \end{aligned}$$

Therefore

$$\begin{aligned} &|\iota_\tau(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u})) x(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \boldsymbol{\Gamma}, \mathbf{H}^{(\tau)}) h(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)| \\ &\leq c_6 I(\mathbf{u} \in U) \prod_{i=1}^m d_i^{\frac{n-m-1}{2}} \prod_{i=m+1}^p d_i^{\frac{n-p-1}{2}} \prod_{j < i \leq m} |d_j - d_i| \prod_{m < j < i} |d_j - d_i| \\ &\times \exp\left(-\frac{1-2a}{2} \operatorname{tr} \sum_{s=1}^2 \mathbf{H}_s^{(\tau)} \mathbf{G}_{ss}(\mathbf{u}) \mathbf{D}_s \mathbf{G}'_{ss}(\mathbf{u}) \mathbf{H}_s^{(\tau)'} \boldsymbol{\Xi}_s^{-1}\right) \\ &\times \operatorname{etr}\left(-\frac{1-2a}{2} \mathbf{Q}_{21} \mathbf{Q}'_{21}\right). \end{aligned} \tag{42}$$

Note that

$$I(\mathbf{u} \in U) \leq I(\mathbf{u} \in C_\epsilon) \leq I(|u_{ij}| = |q_{ij}| < \epsilon, 1 \leq j < i \leq m, m < j < i \leq p).$$

Choose some $\bar{\xi}$ such that $\bar{\xi} > \xi_i$, $i = 1, \dots, p$. Consequently the left-hand side of (42) is bounded by $\bar{h}(\mathbf{d}, \mathbf{q})$, where

$$\begin{aligned}\bar{h}(\mathbf{d}, \mathbf{q}) &= c_6 I(|q_{ij}| < \epsilon, 1 \leq j < i \leq m, m < j < i \leq p) \\ &\times \prod_{i=1}^m d_i^{\frac{n-m-1}{2}} \prod_{i=m+1}^p d_i^{\frac{n-p-1}{2}} \prod_{j < i \leq m} |d_j - d_i| \prod_{m < j < i} |d_j - d_i| \\ &\times \exp\left(-\frac{\bar{\xi}^{-1}}{2}(1-2a) \sum_{i=1}^p d_i\right) \times \text{etr}\left(-\frac{1-2a}{2} \mathbf{Q}_{21} \mathbf{Q}'_{21}\right).\end{aligned}$$

Let $\nu_1 = m(m-1)/2$, $\nu_2 = (p-m)(p-m-1)/2$, $\nu_3 = m(p-m)$. We have

$$\begin{aligned}\int_{R_+^p} \int_{R^{p(p-1)/2}} \bar{h}(\mathbf{d}, \mathbf{q}) d\mathbf{q} d\mathbf{d} &= \int_{R_+^p} \int_{R^{\nu_3}} \int_{R^{\nu_2}} \int_{R^{\nu_1}} \bar{h}(\mathbf{d}, \mathbf{q}) d\mathbf{q}_{11} d\mathbf{q}_{22} d\mathbf{q}_{21} d\mathbf{d} \\ &= c_6 \int_{R_+^p} \prod_{i=1}^m d_i^{\frac{n-m-1}{2}} \prod_{i=m+1}^p d_i^{\frac{n-p-1}{2}} \prod_{j < i \leq m} |d_j - d_i| \prod_{m < j < i} |d_j - d_i| \\ &\times \exp\left(-\frac{\bar{\xi}^{-1}}{2}(1-2a) \sum_{i=1}^p d_i\right) d\mathbf{d} \times \int_{R^{\nu_3}} \text{etr}\left(-\frac{1-2a}{2} \mathbf{Q}_{21} \mathbf{Q}'_{21}\right) d\mathbf{q}_{21} \\ &\times \int_{(-\epsilon, \epsilon)^{\nu_1}} 1 d\mathbf{q}_{11} \int_{(-\epsilon, \epsilon)^{\nu_2}} 1 d\mathbf{q}_{22} < \infty.\end{aligned}$$

The integrability of $\bar{h}(\mathbf{d}, \mathbf{q})$ guarantees the use of the dominated convergence theorem; From (18) and (20)

$$\begin{aligned}\lim_{\beta/\alpha \rightarrow 0} I_\tau &= c_5 \int_{R^{p(p-1)/2}} \int_{R_+^p} \lim_{\beta/\alpha \rightarrow 0} \iota_\tau(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u})) \lim_{\beta/\alpha \rightarrow 0} x(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \Gamma, \mathbf{H}^{(\tau)}) \\ &\times \lim_{\beta/\alpha \rightarrow 0} h(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta) d\mathbf{d} d\mathbf{q} \\ &= c_5 \int_{R^{p(p-1)/2}} \int_{R_+^p} \iota_\tau(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0})) \bar{x}_\Gamma(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}), \mathbf{d}, \mathbf{Q}_{21}, \boldsymbol{\xi}) \\ &\times \lim_{\beta/\alpha \rightarrow 0} h(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta) d\mathbf{d} d\mathbf{q}.\end{aligned}$$

We consider $\lim_{\beta/\alpha \rightarrow 0} h(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)$. First notice that

$$\begin{aligned}\lim_{\beta/\alpha \rightarrow 0} I(\mathbf{d}_1 \in \mathcal{D}_1, \mathbf{d}_2 \in \mathcal{D}_2, (\mathbf{d}_1, \mathbf{d}_2) \in \mathcal{D}_3) &= I(\mathbf{d}_1 \in \mathcal{D}_1) I(\mathbf{d}_2 \in \mathcal{D}_2), \\ \lim_{\beta/\alpha \rightarrow 0} \prod_{j \leq m < i} \left(1 - \frac{\beta d_i}{\alpha d_j}\right) &= 1.\end{aligned}$$

From (17), we find

$$\begin{aligned}\lim_{\beta/\alpha \rightarrow 0} J^*(\mathbf{u}) &= J^*(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}), \\ \lim_{\beta/\alpha \rightarrow 0} I(\mathbf{u} \in U) &= I((\mathbf{q}_{11}, \mathbf{q}_{22}) = (\mathbf{u}_{11}, \mathbf{u}_{22}) \in U_0),\end{aligned}$$

where $U_0 = \{(\mathbf{u}_{11}, \mathbf{u}_{22}) | (\mathbf{u}_{11}, \mathbf{u}_{22}, \mathbf{0}) \in U\}$ denotes the slice of U by $\mathbf{u}_{12} = 0$, and that

$$\lim_{\beta/\alpha \rightarrow 0} \mathbf{G}_{11}(\mathbf{u}) = \mathbf{G}_{11}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}) \in \mathcal{O}^+(m),$$

$$\begin{aligned}
\lim_{\beta/\alpha \rightarrow 0} \mathbf{G}_{22}(\mathbf{u}) &= \mathbf{G}_{22}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}) \in \mathcal{O}^+(p-m), \\
\lim_{\beta/\alpha \rightarrow 0} \mathbf{G}_{21}(\mathbf{u}) &= \mathbf{0}, \\
\lim_{\beta/\alpha \rightarrow 0} \text{etr}\left(-\frac{\beta}{2\alpha} \mathbf{H}_1^{(\tau)} \mathbf{G}_{12}(\mathbf{u}) \mathbf{D}_2 \mathbf{G}'_{12}(\mathbf{u}) \mathbf{H}_1^{(\tau)'} \boldsymbol{\Xi}_1^{-1}\right) &= 1.
\end{aligned}$$

Since $d\mu$ is invariant, especially w.r.t. both of the transformations

$$\mathbf{G} \rightarrow \text{diag}(\mathbf{H}_1, \mathbf{H}_2)\mathbf{G}, \quad \mathbf{G} \rightarrow \mathbf{G} \text{diag}(\mathbf{H}_1, \mathbf{H}_2), \quad (43)$$

the measure on U_0 given by

$$J^*(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}) d\mathbf{q}_{11} d\mathbf{q}_{22} \quad (44)$$

induces the invariant measure on V_0 , the slice of V by $\mathbf{G}_{12} = \mathbf{0}$, w.r.t. (43) through

$$\mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}) = \text{diag}(\mathbf{G}_{11}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}), \mathbf{G}_{22}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0})). \quad (45)$$

If \mathbf{G}_{11} and \mathbf{G}_{22} independently follow the invariant probability distributions respectively on $\mathcal{O}^+(m)$ and $\mathcal{O}^+(p-m)$, then the distribution on V_0 given by

$$\mathbf{G}_0 = \text{diag}(\mathbf{G}_{11}, \mathbf{G}_{22}) \quad (46)$$

is also invariant w.r.t. the transformations (43), hence must be proportional to the above-mentioned distribution on V_0 given by (45) and (44). Consequently

$$\begin{aligned}
\lim_{\beta/\alpha \rightarrow 0} I_\tau &= c_6 \int_{R^{m(p-m)}} \int_{R_+^p} \int_{V_0} \iota_\tau(\mathbf{H}^{(\tau)} \mathbf{G}_0) \bar{x}_\Gamma(\mathbf{H}^{(\tau)} \mathbf{G}_0, \mathbf{d}, \mathbf{Q}_{21}, \boldsymbol{\xi}) I(\mathbf{d}_1 \in \mathcal{D}_1) I(\mathbf{d}_2 \in \mathcal{D}_2) \\
&\quad \times \prod_{i=1}^m d_i^{\frac{n-m-1}{2}} \prod_{i=m+1}^p d_i^{\frac{n-p-1}{2}} \prod_{j<i \leq m} (d_j - d_i) \prod_{m<j<i} (d_j - d_i) \\
&\quad \times \exp\left(-\frac{1}{2} \text{tr} \sum_{s=1}^2 \mathbf{H}_s^{(\tau)} \mathbf{G}_{ss} \mathbf{D}_s \mathbf{G}'_{ss} \mathbf{H}_s^{(\tau)'} \boldsymbol{\Xi}_s^{-1}\right) \\
&\quad \times \text{etr}\left(-\frac{1}{2} \mathbf{Q}_{21} \mathbf{Q}'_{21}\right) d\mu_1(\mathbf{G}_{11}) d\mu_2(\mathbf{G}_{22}) d\mathbf{d} d\mathbf{q}_{21},
\end{aligned}$$

where \mathbf{G}_0 is given by (46), and μ_1, μ_2 are the invariant probability measures respectively on $\mathcal{O}^+(m)$ and $\mathcal{O}^+(p-m)$.

Let $O_0^{(\tau)}$ denote the slice of $O^{(\tau)}$ by $\mathbf{G}_{12} = \mathbf{0}$. Since $O^{(\tau)} = \mathbf{H}^{(\tau)} V$, $O_0^{(\tau)} = \mathbf{H}^{(\tau)} V_0$. Consequently for each $1 \leq \tau \leq T$,

$$\begin{aligned}
\lim_{\beta/\alpha \rightarrow 0} I_\tau &= c_6 \int_{R^{m(p-m)}} \int_{R_+^p} \int_{O_0^{(\tau)}} \iota_\tau(\mathbf{G}_0) \bar{x}_\Gamma(\mathbf{G}_0, \mathbf{d}, \mathbf{Q}_{21}, \boldsymbol{\xi}) I(\mathbf{d}_1 \in \mathcal{D}_1) I(\mathbf{d}_2 \in \mathcal{D}_2) \\
&\quad \times \prod_{i=1}^m d_i^{\frac{n-m-1}{2}} \prod_{i=m+1}^p d_i^{\frac{n-p-1}{2}} \prod_{j<i \leq m} (d_j - d_i) \prod_{m<j<i} (d_j - d_i) \\
&\quad \times \exp\left(-\frac{1}{2} \text{tr} \sum_{s=1}^2 \mathbf{G}_{ss} \mathbf{D}_s \mathbf{G}'_{ss} \boldsymbol{\Xi}_s^{-1}\right) \text{etr}\left(-\frac{1}{2} \mathbf{Q}_{21} \mathbf{Q}'_{21}\right) d\mu_1(\mathbf{G}_{11}) d\mu_2(\mathbf{G}_{22}) d\mathbf{d} d\mathbf{q}_{21}.
\end{aligned}$$

Note that $\bigcup_{\tau=1}^T O_0^{(\tau)} = \mathcal{O}(m, p-m)$ and $\iota_\tau(\mathbf{G}_0)$ vanishes on $\mathcal{O}(m, p-m) \setminus O_0^{(\tau)}$. Therefore we have

$$\begin{aligned} \lim_{\beta/\alpha \rightarrow 0} I_\tau &= c_6 \int_{R^{m(p-m)}} \int_{R_+^p} \int_{\mathcal{O}(m, p-m)} \iota_\tau(\mathbf{G}_0) \bar{x}_\Gamma(\mathbf{G}_0, \mathbf{d}, \mathbf{Q}_{21}, \boldsymbol{\xi}) I(\mathbf{d}_1 \in \mathcal{D}_1) I(\mathbf{d}_2 \in \mathcal{D}_2) \\ &\quad \times \prod_{i=1}^m d_i^{\frac{n-m-1}{2}} \prod_{i=m+1}^p d_i^{\frac{n-p-1}{2}} \prod_{j < i \leq m} (d_j - d_i) \prod_{m < j < i} (d_j - d_i) \\ &\quad \times \exp\left(-\frac{1}{2} \operatorname{tr} \sum_{s=1}^2 \mathbf{G}_{ss} \mathbf{D}_s \mathbf{G}'_{ss} \boldsymbol{\Xi}_s^{-1}\right) \operatorname{etr}\left(-\frac{1}{2} \mathbf{Q}_{21} \mathbf{Q}'_{21}\right) d\mu_1(\mathbf{G}_{11}) d\mu_2(\mathbf{G}_{22}) d\mathbf{d} d\mathbf{q}_{21}. \end{aligned} \quad (47)$$

From (35), (37) and (47), we have

$$\begin{aligned} &\lim_{\beta/\alpha \rightarrow 0} E[x(\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta)] \\ &= c_6 \int_{R^{m(p-m)}} \int_{R_+^p} \int_{\mathcal{O}^+(p-m)} \int_{\mathcal{O}^+(m)} \bar{x}_\Gamma(\mathbf{G}_0, \mathbf{d}, \mathbf{Q}_{21}, \boldsymbol{\xi}) I(\mathbf{d}_1 \in \mathcal{D}_1) I(\mathbf{d}_2 \in \mathcal{D}_2) \\ &\quad \times \prod_{i=1}^m d_i^{\frac{n-m-1}{2}} \prod_{i=m+1}^p d_i^{\frac{n-p-1}{2}} \prod_{j < i \leq m} (d_j - d_i) \prod_{m < j < i} (d_j - d_i) \\ &\quad \times \exp\left(-\frac{1}{2} \operatorname{tr} \sum_{s=1}^2 \mathbf{G}_{ss} \mathbf{D}_s \mathbf{G}'_{ss} \boldsymbol{\Xi}_s^{-1}\right) \times \operatorname{etr}\left(-\frac{1}{2} \mathbf{Q}_{21} \mathbf{Q}'_{21}\right) d\mu_1(\mathbf{G}_{11}) d\mu_2(\mathbf{G}_{22}) d\mathbf{d} d\mathbf{q}_{21}. \end{aligned} \quad (48)$$

Under the distribution (22) and the spectral decompositions (23), the joint density function of $(\mathbf{d}_1, \mathbf{G}_{11})$ ($(\mathbf{d}_2, \mathbf{G}_{22})$) with respect to the product measure of Lebesgue measure on R_+^m (R_+^{p-m}) and the invariant probability measure μ_1 (μ_2) on $\mathcal{O}^+(m)$ ($\mathcal{O}^+(p-m)$) is given by the following functions, $F_1(\mathbf{d}_1, \mathbf{G}_{11})$ ($F_2(\mathbf{d}_2, \mathbf{G}_{22})$);

$$\begin{aligned} F_1(\mathbf{G}_{11}, \mathbf{d}_1) &= K_1 |\boldsymbol{\Xi}_1|^{-\frac{n}{2}} \prod_{i=1}^m d_i^{\frac{n-m-1}{2}} \prod_{1 \leq j < i \leq m} (d_j - d_i) \operatorname{etr}\left(-\frac{1}{2} \mathbf{G}_{11} \mathbf{D}_1 \mathbf{G}'_{11} \boldsymbol{\Xi}_1^{-1}\right) \\ F_2(\mathbf{G}_{22}, \mathbf{d}_2) &= K_2 |\boldsymbol{\Xi}_2|^{-\frac{n-m}{2}} \prod_{i=m+1}^p d_i^{\frac{n-p-1}{2}} \prod_{m < j < i \leq p} (d_j - d_i) \operatorname{etr}\left(-\frac{1}{2} \mathbf{G}_{22} \mathbf{D}_2 \mathbf{G}'_{22} \boldsymbol{\Xi}_2^{-1}\right), \end{aligned}$$

with K_1, K_2 as normalizing constants. The density function of \mathbf{Z}_{21} is given by

$$F_3(\mathbf{z}_{21}) = K_3 \operatorname{etr}\left(-\frac{1}{2} \mathbf{Z}_{21} \mathbf{Z}'_{21}\right),$$

where K_3 is a normalizing constant. Using $F_1(\mathbf{G}_{11}, \mathbf{d}_1)$, $F_2(\mathbf{G}_{22}, \mathbf{d}_2)$, $F_3(\mathbf{z}_{21})$, we can rewrite the right-hand side of (48) as

$$\begin{aligned} &c_7 \int_{R^{m(p-m)}} \int_{\mathcal{D}_2} \int_{\mathcal{D}_1} \int_{\mathcal{O}^+(p-m)} \int_{\mathcal{O}^+(m)} \bar{x}_\Gamma(\mathbf{G}_0, (\mathbf{d}_1, \mathbf{d}_2), \mathbf{Z}_{21}, \boldsymbol{\xi}) \\ &\quad \times F_1(\mathbf{G}_{11}, \mathbf{d}_1) F_2(\mathbf{G}_{22}, \mathbf{d}_2) F_3(\mathbf{z}_{21}) d\mu_1(\mathbf{G}_{11}) d\mu_2(\mathbf{G}_{22}) d\mathbf{d}_1 d\mathbf{d}_2 d\mathbf{z}_{21}. \end{aligned}$$

If we consider the special case $x(\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) = 1$, we notice that $c_7 = 1$.

A.2 Proof of Lemma 2

Using Lemma 1, we will calculate

$$\begin{aligned} & \lim_{\beta/\alpha \rightarrow 0} E[\text{tr} \mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \mathbf{\Gamma} \mathbf{\Lambda}^{-1} \mathbf{\Gamma}'], \\ & \lim_{\beta/\alpha \rightarrow 0} E[\text{tr} (\mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \mathbf{\Gamma} \mathbf{\Lambda}^{-1} \mathbf{\Gamma}')^2]. \end{aligned}$$

Let

$$\begin{aligned} x_1(\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) &= \text{tr}(\mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \mathbf{\Gamma} \mathbf{\Lambda}^{-1} \mathbf{\Gamma}'), \\ x_2(\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) &= \text{tr}(\mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \mathbf{\Gamma} \mathbf{\Lambda}^{-1} \mathbf{\Gamma}')^2, \end{aligned}$$

then

$$\begin{aligned} x_1(\mathbf{\Gamma} \mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) &= \sum_{i=1}^p \sum_{j=1}^p \lambda_i^{-1} l_j c_j g_{ij}^2 \\ &\leq (\max_j c_j) \sum_{i=1}^p \sum_{j=1}^p \lambda_i^{-1} l_j g_{ij}^2 = 3(\max_j c_j) \text{tr} \left(\frac{1}{3} \mathbf{GLG}' \mathbf{\Lambda}^{-1} \right) \\ &\leq 3(\max_j c_j) \text{etr} \left(\frac{1}{3} \mathbf{GLG}' \mathbf{\Lambda}^{-1} \right), \\ x_2(\mathbf{\Gamma} \mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) &= \text{tr}(\mathbf{\Lambda}^{-1/2} \mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \mathbf{\Lambda}^{-1/2})^2 \\ &\leq \left(\text{tr} \mathbf{\Lambda}^{-1/2} \mathbf{GL}^{1/2} \mathbf{CL}^{1/2} \mathbf{G}' \mathbf{\Lambda}^{-1/2} \right)^2 = \left(\sum_{i=1}^p \sum_{j=1}^p \lambda_i^{-1} l_j c_j g_{ij}^2 \right)^2 \\ &\leq \left((\max_j c_j) \sum_{i=1}^p \sum_{j=1}^p \lambda_i^{-1} l_j g_{ij}^2 \right)^2 = \left\{ 6(\max_j c_j) \text{tr} \left(\frac{1}{6} \mathbf{GLG}' \mathbf{\Lambda}^{-1} \right) \right\}^2 \\ &\leq \left\{ 6(\max_j c_j) \text{etr} \left(\frac{1}{6} \mathbf{GLG}' \mathbf{\Lambda}^{-1} \right) \right\}^2 = 36(\max_j c_j)^2 \text{etr} \left(\frac{1}{3} \mathbf{GLG}' \mathbf{\Lambda}^{-1} \right), \end{aligned}$$

hence (19) is satisfied for both x_1 and x_2 . Now let

$$\mathbf{B}(\mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}) = \mathbf{\Lambda}^{-1/2} \mathbf{H}^{(\tau)} \mathbf{GL}^{1/2}$$

for each τ . Then we have

$$\begin{aligned} x_1(\mathbf{\Gamma} \mathbf{H}^{(\tau)} \mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) &= \text{tr} \mathbf{BCB}', \\ x_2(\mathbf{\Gamma} \mathbf{H}^{(\tau)} \mathbf{G}, \mathbf{l}, \boldsymbol{\lambda}, \alpha, \beta) &= \text{tr} (\mathbf{BCB}')^2. \end{aligned}$$

We notice that

$$\begin{aligned} \mathbf{B} &= \mathbf{\Lambda}^{-1/2} \mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{u}) \mathbf{L}^{1/2} \\ &= \begin{pmatrix} \mathbf{\Lambda}_1^{-1/2} \mathbf{H}_1^{(\tau)} \mathbf{G}_{11}(\mathbf{u}) \mathbf{L}_1^{1/2} & \mathbf{\Lambda}_1^{-1/2} \mathbf{H}_1^{(\tau)} \mathbf{G}_{12}(\mathbf{u}) \mathbf{L}_2^{1/2} \\ \mathbf{\Lambda}_2^{-1/2} \mathbf{H}_2^{(\tau)} \mathbf{U}_{21} \mathbf{L}_1^{1/2} & \mathbf{\Lambda}_2^{-1/2} \mathbf{H}_2^{(\tau)} \mathbf{G}_{22}(\mathbf{u}) \mathbf{L}_2^{1/2} \end{pmatrix} \\ &= \begin{pmatrix} \boldsymbol{\Xi}_1^{-1/2} \mathbf{H}_1^{(\tau)} \mathbf{G}_{11}(\mathbf{u}) \mathbf{D}_1^{1/2} & \alpha^{-1/2} \beta^{1/2} \boldsymbol{\Xi}_1^{-1/2} \mathbf{H}_1^{(\tau)} \mathbf{G}_{12}(\mathbf{u}) \mathbf{D}_2^{1/2} \\ \alpha^{1/2} \beta^{-1/2} \boldsymbol{\Xi}_2^{-1/2} \mathbf{H}_2^{(\tau)} \mathbf{U}_{21} \mathbf{D}_1^{1/2} & \boldsymbol{\Xi}_2^{-1/2} \mathbf{H}_2^{(\tau)} \mathbf{G}_{22}(\mathbf{u}) \mathbf{D}_2^{1/2} \end{pmatrix} \\ &= \begin{pmatrix} \boldsymbol{\Xi}_1^{-1/2} \mathbf{H}_1^{(\tau)} \mathbf{G}_{11}(\mathbf{u}) \mathbf{D}_1^{1/2} & \alpha^{-1/2} \beta^{1/2} \boldsymbol{\Xi}_1^{-1/2} \mathbf{H}_1^{(\tau)} \mathbf{G}_{12}(\mathbf{u}) \mathbf{D}_2^{1/2} \\ \mathbf{Q}_{21} & \boldsymbol{\Xi}_2^{-1/2} \mathbf{H}_2^{(\tau)} \mathbf{G}_{22}(\mathbf{u}) \mathbf{D}_2^{1/2} \end{pmatrix}. \end{aligned}$$

Substitute $\mathbf{u}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)$ with \mathbf{u} in the last matrix and denote it by $\mathbf{B}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)$. Then

$$\begin{aligned} x_1(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \boldsymbol{\Gamma}, \mathbf{H}^{(\tau)}) &= \text{tr} \mathbf{B}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta) \mathbf{C} \mathbf{B}'(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta), \\ x_2(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \boldsymbol{\Gamma}, \mathbf{H}^{(\tau)}) &= \text{tr}(\mathbf{B}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta) \mathbf{C} \mathbf{B}'(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta))^2. \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{\beta/\alpha \rightarrow 0} x_1(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \boldsymbol{\Gamma}, \mathbf{H}^{(\tau)}) &= \text{tr} \bar{\mathbf{B}} \mathbf{C} \bar{\mathbf{B}}', \\ \lim_{\beta/\alpha \rightarrow 0} x_2(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta; \boldsymbol{\Gamma}, \mathbf{H}^{(\tau)}) &= \text{tr}(\bar{\mathbf{B}} \mathbf{C} \bar{\mathbf{B}}')^2, \end{aligned}$$

where

$$\bar{\mathbf{B}} = \begin{pmatrix} \bar{\mathbf{B}}_{11} & \bar{\mathbf{B}}_{12} \\ \bar{\mathbf{B}}_{21} & \bar{\mathbf{B}}_{22} \end{pmatrix} = \lim_{\beta/\alpha \rightarrow 0} \mathbf{B}(\mathbf{d}, \mathbf{q}, \boldsymbol{\xi}, \alpha, \beta)$$

is given by

$$\begin{aligned} \bar{\mathbf{B}}_{11} &= \boldsymbol{\Xi}_1^{-1/2} \mathbf{H}_1^{(\tau)} \mathbf{G}_{11}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}) \mathbf{D}_1^{1/2} = \boldsymbol{\Xi}_1^{-1/2} (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))_{11} \mathbf{D}_1^{1/2}, \\ \bar{\mathbf{B}}_{12} &= \mathbf{0}, \\ \bar{\mathbf{B}}_{21} &= \mathbf{Q}_{21}, \\ \bar{\mathbf{B}}_{22} &= \boldsymbol{\Xi}_2^{-1/2} \mathbf{H}_2^{(\tau)} \mathbf{G}_{22}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}) \mathbf{D}_2^{1/2} = \boldsymbol{\Xi}_2^{-1/2} (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))_{22} \mathbf{D}_2^{1/2}, \end{aligned}$$

because of (18). By straightforward calculation we have

$$\begin{aligned} \text{tr}(\bar{\mathbf{B}} \mathbf{C} \bar{\mathbf{B}}') &= \text{tr} \bar{\mathbf{B}}_{11} \mathbf{C}_1 \bar{\mathbf{B}}'_{11} + \text{tr} \bar{\mathbf{B}}_{12} \mathbf{C}_2 \bar{\mathbf{B}}'_{12} + \text{tr} \bar{\mathbf{B}}_{21} \mathbf{C}_1 \bar{\mathbf{B}}'_{21} + \text{tr} \bar{\mathbf{B}}_{22} \mathbf{C}_2 \bar{\mathbf{B}}'_{22} \\ &= \sum_{s=1}^2 \text{tr}(\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))_{ss} \mathbf{D}_s^{1/2} \mathbf{C}_s \mathbf{D}_s^{1/2} (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))'_{ss} \boldsymbol{\Xi}_s^{-1} + \text{tr} \mathbf{Q}_{21} \mathbf{C}_1 \mathbf{Q}'_{21}, \end{aligned}$$

$$\begin{aligned} \text{tr}(\bar{\mathbf{B}} \mathbf{C} \bar{\mathbf{B}}')^2 &= \text{tr}(\mathbf{C} \bar{\mathbf{B}}' \bar{\mathbf{B}})^2 \\ &= \text{tr} \begin{pmatrix} \mathbf{C}_1 (\bar{\mathbf{B}}'_{11} \bar{\mathbf{B}}_{11} + \bar{\mathbf{B}}'_{21} \bar{\mathbf{B}}_{21}) & \mathbf{C}_1 \bar{\mathbf{B}}'_{21} \bar{\mathbf{B}}_{22} \\ \mathbf{C}_2 \bar{\mathbf{B}}'_{22} \bar{\mathbf{B}}_{21} & \mathbf{C}_2 \bar{\mathbf{B}}'_{22} \bar{\mathbf{B}}_{22} \end{pmatrix}^2 \\ &= \text{tr}(\mathbf{C}_1 (\bar{\mathbf{B}}'_{11} \bar{\mathbf{B}}_{11} + \bar{\mathbf{B}}'_{21} \bar{\mathbf{B}}_{21}))^2 + 2 \text{tr} \mathbf{C}_1 \bar{\mathbf{B}}'_{21} \bar{\mathbf{B}}_{22} \mathbf{C}_2 \bar{\mathbf{B}}'_{22} \bar{\mathbf{B}}_{21} + \text{tr}(\mathbf{C}_2 \bar{\mathbf{B}}'_{22} \bar{\mathbf{B}}_{22})^2 \\ &= \text{tr}(\mathbf{C}_1 \mathbf{D}_1^{1/2} (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))'_{11} \boldsymbol{\Xi}_1^{-1} (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))_{11} \mathbf{D}_1^{1/2} + \mathbf{C}_1 \mathbf{Q}'_{21} \mathbf{Q}_{21})^2 \\ &\quad + 2 \text{tr}(\mathbf{C}_1 \mathbf{Q}'_{21} \boldsymbol{\Xi}_2^{-1/2} (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))_{22} \mathbf{D}_2^{1/2} \mathbf{C}_2 \\ &\quad \quad \times \mathbf{D}_2^{1/2} (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))'_{22} \boldsymbol{\Xi}_2^{-1/2} \mathbf{Q}_{21}) \\ &\quad + \text{tr}(\mathbf{C}_2 \mathbf{D}_2^{1/2} (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))'_{22} \boldsymbol{\Xi}_2^{-1} (\mathbf{H}^{(\tau)} \mathbf{G}(\mathbf{q}_{11}, \mathbf{q}_{22}, \mathbf{0}))_{22} \mathbf{D}_2^{1/2})^2. \end{aligned}$$

Consequently we have the following results; all the asymptotic expectations below are taken with respect to the distributions in (22) and the spectral decompositions (23).

$$\begin{aligned} &\lim_{\beta/\alpha \rightarrow 0} E[\text{tr} \mathbf{G} \mathbf{L}^{1/2} \mathbf{C} \mathbf{L}^{1/2} \mathbf{G}' \boldsymbol{\Gamma} \boldsymbol{\Lambda}^{-1} \boldsymbol{\Gamma}'] \\ &= E[\text{tr} \mathbf{G}_{11} \mathbf{D}_1^{1/2} \mathbf{C}_1 \mathbf{D}_1^{1/2} \mathbf{G}'_{11} \boldsymbol{\Xi}_1^{-1}] + E[\text{tr} \mathbf{G}_{22} \mathbf{D}_2^{1/2} \mathbf{C}_2 \mathbf{D}_2^{1/2} \mathbf{G}'_{22} \boldsymbol{\Xi}_2^{-1}] \\ &\quad + E[\text{tr} \mathbf{Z}_{21} \mathbf{C}_1 \mathbf{Z}'_{21}] \end{aligned} \quad (49)$$

$$\begin{aligned}
& \lim_{\beta/\alpha \rightarrow 0} E[\text{tr}(\mathbf{G}\mathbf{L}^{1/2}\mathbf{C}\mathbf{L}^{1/2}\mathbf{G}'\mathbf{\Gamma}\mathbf{\Lambda}^{-1}\mathbf{\Gamma}')^2] \\
&= E[\text{tr}(\mathbf{C}_1\mathbf{D}_1^{1/2}\mathbf{G}'_{11}\mathbf{\Xi}_1^{-1}\mathbf{G}_{11}\mathbf{D}_1^{1/2} + \mathbf{C}_1\mathbf{Z}'_{21}\mathbf{Z}_{21})^2] \\
&\quad + 2E[\text{tr}\mathbf{C}_1\mathbf{Z}'_{21}\mathbf{\Xi}_2^{-1/2}\mathbf{G}_{22}\mathbf{D}_2^{1/2}\mathbf{C}_2\mathbf{D}_2^{1/2}\mathbf{G}'_{22}\mathbf{\Xi}_2^{-1/2}\mathbf{Z}_{21}] \\
&\quad + E[\text{tr}(\mathbf{C}_2\mathbf{D}_2^{1/2}\mathbf{G}'_{22}\mathbf{\Xi}_2^{-1}\mathbf{G}_{22}\mathbf{D}_2^{1/2})^2] \\
&= E[\text{tr}(\mathbf{G}_{11}\mathbf{D}_1^{1/2}\mathbf{C}_1\mathbf{D}_1^{1/2}\mathbf{G}'_{11}\mathbf{\Xi}_1^{-1})^2] \\
&\quad + 2\text{tr} E[\mathbf{C}_1\mathbf{D}_1^{1/2}\mathbf{G}'_{11}\mathbf{\Xi}_1^{-1}\mathbf{G}_{11}\mathbf{D}_1^{1/2}\mathbf{C}_1]E[\mathbf{Z}'_{21}\mathbf{Z}_{21}] \\
&\quad + E[\text{tr}\mathbf{C}_1\mathbf{Z}'_{21}\mathbf{Z}_{21}\mathbf{C}_1\mathbf{Z}'_{21}\mathbf{Z}_{21}] \\
&\quad + 2\text{tr} E[\mathbf{\Xi}_2^{-1/2}\mathbf{G}_{22}\mathbf{D}_2^{1/2}\mathbf{C}_2\mathbf{D}_2^{1/2}\mathbf{G}'_{22}\mathbf{\Xi}_2^{-1/2}]E[\mathbf{Z}_{21}\mathbf{C}_1\mathbf{Z}'_{21}] \\
&\quad + E[\text{tr}(\mathbf{G}_{22}\mathbf{D}_2^{1/2}\mathbf{C}_2\mathbf{D}_2^{1/2}\mathbf{G}'_{22}\mathbf{\Xi}_2^{-1})^2].
\end{aligned} \tag{50}$$

We further calculate the expectations related to \mathbf{Z}_{21} . It is obvious that

$$E[\mathbf{Z}_{21}\mathbf{C}_1\mathbf{Z}'_{21}] = (\text{tr}\mathbf{C}_1)\mathbf{I}_{p-m}, \quad E[\mathbf{Z}'_{21}\mathbf{Z}_{21}] = (p-m)\mathbf{I}_m, \tag{51}$$

since $(\mathbf{Z}_{21})_{ij}$, $1 \leq i \leq p-m$, $1 \leq j \leq m$, are all independently distributed as the standard normal distributions. Letting $\mathbf{T} = (t_{ij}) = \mathbf{Z}'_{21}\mathbf{Z}_{21}$ we have

$$\begin{aligned}
E[\text{tr}\mathbf{C}_1\mathbf{Z}'_{21}\mathbf{Z}_{21}\mathbf{C}_1\mathbf{Z}'_{21}\mathbf{Z}_{21}] &= E[\text{tr}\mathbf{C}_1\mathbf{T}\mathbf{C}_1\mathbf{T}] = \sum_{i=1}^m c_i E[(\mathbf{T}\mathbf{C}_1\mathbf{T})_{ii}] \\
&= \sum_{i=1}^m c_i E[\sum_{s=1}^m t_{is}^2 c_s] = \sum_{i=1}^m \sum_{s=1}^m c_i c_s E[t_{is}^2],
\end{aligned} \tag{52}$$

while

$$\begin{aligned}
E[t_{is}^2] &= E[(\sum_{j=1}^{p-m} (\mathbf{Z}_{21})_{ji}(\mathbf{Z}_{21})_{js})^2] \\
&= E[\sum_{j=1}^{p-m} (\mathbf{Z}_{21})_{ji}^2 (\mathbf{Z}_{21})_{js}^2 + 2 \sum_{j_1 < j_2} (\mathbf{Z}_{21})_{j_1 i} (\mathbf{Z}_{21})_{j_1 s} (\mathbf{Z}_{21})_{j_2 i} (\mathbf{Z}_{21})_{j_2 s}].
\end{aligned} \tag{53}$$

We also have

$$E[(\mathbf{Z}_{21})_{ji}^2 (\mathbf{Z}_{21})_{js}^2] = \begin{cases} 3, & \text{if } i = s, \\ 1, & \text{if } i \neq s, \end{cases} \tag{54}$$

$$E[(\mathbf{Z}_{21})_{j_1 i} (\mathbf{Z}_{21})_{j_1 s} (\mathbf{Z}_{21})_{j_2 i} (\mathbf{Z}_{21})_{j_2 s}] = \begin{cases} 1, & \text{if } i = s, \\ 0, & \text{if } i \neq s. \end{cases} \tag{55}$$

Substituting (54),(55) into (53), we have

$$E[t_{is}^2] = \begin{cases} (p-m)(p-m+2), & \text{if } i = s, \\ p-m, & \text{if } i \neq s. \end{cases} \tag{56}$$

Consequently from (52) and (56),

$$E[\text{tr}\mathbf{C}_1\mathbf{Z}'_{21}\mathbf{Z}_{21}\mathbf{C}_1\mathbf{Z}'_{21}\mathbf{Z}_{21}] = (p-m)(p-m+2) \sum_{i=1}^m c_i^2 + 2(p-m) \sum_{1 \leq i < s \leq m} c_i c_s. \tag{57}$$

Substituting (51) and (57) into (49), (50), we have the result.

A.3 Analytic Evaluation of Asymptotic Risk

We illustrate an analytic calculation of $E[d_i]$, $E[d_i^2]$, $i = 1, \dots, p$, by the case $p = 4$, $m = 1$ and $n (\geq 4)$ even. Suppose $\mathbf{S} \sim \mathbf{W}_3(n, \mathbf{I}_p)$. Note the density function of $\mathbf{l} = (l_1, l_2, l_3)$ is given by (see e.g. Theorem 3.2.18 of Muirhead (1982))

$$K_3(n) \prod_{i=1}^3 l_i^u (l_1 - l_2)(l_1 - l_3)(l_2 - l_3) \exp\left(-\frac{1}{2} \sum_{i=1}^3 l_i\right),$$

where $u = u(n) = (n - 4)/2$, which is an integer, and

$$K_3(n) = \pi^{3/2} / \left(2^{3n/2} \Gamma(n/2) \Gamma((n-1)/2) \Gamma(n/2 - 1) \Gamma(3/2) \Gamma(1) \Gamma(1/2)\right).$$

Let

$$\Delta_1 = l_1 - l_2, \quad \Delta_2 = l_2 - l_3, \quad \Delta_3 = l_3.$$

The density function $f_3(\mathbf{\Delta})$ of $\mathbf{\Delta} = (\Delta_1, \Delta_2, \Delta_3)$ is given by

$$\begin{aligned} f_3(\mathbf{\Delta}) &= K_3(n) \Delta_3^u (\Delta_2 + \Delta_3)^u (\Delta_1 + \Delta_2 + \Delta_3)^u \\ &\quad \times \Delta_1 \Delta_2 (\Delta_1 + \Delta_2) \exp\left(-\frac{1}{2} (\Delta_1 + 2\Delta_2 + 3\Delta_3)\right) \\ &= K_3(n) \left(\sum_{i=0}^u \binom{u}{i} \Delta_2^i \Delta_3^{u-i}\right) \left(\sum_{s=0}^u \sum_{t=0}^{u-s} \binom{u}{s} \binom{u-s}{t} \Delta_1^s \Delta_2^t \Delta_3^{u-s-t}\right) \\ &\quad \times \left(\sum_{j=0}^1 \Delta_1^j \Delta_2^{1-j}\right) \Delta_1 \Delta_2 \Delta_3^u \exp\left(-\frac{1}{2} (\Delta_1 + 2\Delta_2 + 3\Delta_3)\right) \\ &= K_3(n) \sum_{i=0}^u \sum_{j=0}^1 \sum_{s=0}^u \sum_{t=0}^{u-s} \binom{u}{i} \binom{u}{s} \binom{u-s}{t} \Delta_1^{j+s+1} \Delta_2^{i-j+t+2} \Delta_3^{3u-i-s-t} \\ &\quad \times \exp\left(-\frac{1}{2} (\Delta_1 + 2\Delta_2 + 3\Delta_3)\right) \end{aligned}$$

We define a function $F_3(x_1, x_2, x_3; n)$ of nonnegative integers x_i , $i = 1, 2, 3$, as

$$F_3(x_1, x_2, x_3; n) = E[\Delta_1^{x_1} \Delta_2^{x_2} \Delta_3^{x_3}].$$

Then

$$\begin{aligned} F_3(x_1, x_2, x_3; n) &= K_3(n) \sum_{i=0}^u \sum_{j=0}^1 \sum_{s=0}^u \sum_{t=0}^{u-s} \binom{u}{i} \binom{u}{s} \binom{u-s}{t} \\ &\quad \times \int_0^\infty \Delta_1^{j+s+x_1+1} \exp\left(-\frac{1}{2} \Delta_1\right) d\Delta_1 \\ &\quad \times \int_0^\infty \Delta_2^{i-j+t+x_2+2} \exp(-\Delta_2) d\Delta_2 \\ &\quad \times \int_0^\infty \Delta_3^{3u-i-s-t+x_3} \exp\left(-\frac{3}{2} \Delta_3\right) d\Delta_3 \\ &= K_3(n) \sum_{i=0}^u \sum_{j=0}^1 \sum_{s=0}^u \sum_{t=0}^{u-s} \binom{u}{i} \binom{u}{s} \binom{u-s}{t} \\ &\quad \times 2^{3u-i+j-t+x_1+x_3+3} 3^{-3u+i+s+t-x_3-1} \\ &\quad \times (j+s+x_1+1)! (i-j+t+2+x_2)! \\ &\quad \times (3u-i-s-t+x_3)! \end{aligned}$$

Note that for the case $p = 4$, $m = 1$, the distributions of d_i , $i = 1, \dots, 4$, in Theorem 5 is given as follows; $d_1 = \mathbf{W}_{11} \sim \chi_n^2$ and $d_2 > d_3 > d_4$ are the ordered eigenvalues of $\mathbf{W}_{22} \sim \mathbf{W}_3(n-1, \mathbf{I}_3)$. Using $\Delta_1 = d_2 - d_3$, $\Delta_2 = d_3 - d_4$, $\Delta_3 = d_4$ and $F_3(x_1, x_2, x_3; n)$ as above, we can calculate $\mathbf{b} = (b_1, \dots, b_4)$ and $\mathbf{A} = (a_{ij})_{1 \leq i, j \leq 4}$ in Theorem 5 as follows;

$$\begin{aligned}
b_1 &= E[d_1] + (p - m) = n + 3, \\
b_2 &= E[d_2] = E[\Delta_1 + \Delta_2 + \Delta_3] = F_3(1, 0, 0; n - 1) + F_3(0, 1, 0; n - 1) \\
&\quad + F_3(0, 0, 1; n - 1), \\
b_3 &= E[d_3] = E[\Delta_2 + \Delta_3] = F_3(0, 1, 0; n - 1) + F_3(0, 0, 1; n - 1), \\
b_4 &= E[d_4] = E[\Delta_3] = F_3(0, 0, 1; n - 1), \\
a_{11} &= E[d_1^2 + 2(p - m)d_1] + (p - m)(p - m + 2) \\
&= n^2 + 2n + 6n + 15 = n^2 + 8n + 15, \\
a_{22} &= E[d_2^2] = E\left[\sum_{i=1}^3 \Delta_i^2 + 2 \sum_{1 \leq i < j \leq 3} \Delta_i \Delta_j\right] \\
&= F_3(2, 0, 0; n - 1) + F_3(0, 2, 0; n - 1) + F_3(0, 0, 2; n - 1) \\
&\quad + 2F_3(1, 1, 0; n - 1) + 2F_3(1, 0, 1; n - 1) + 2F_3(0, 1, 1; n - 1), \\
a_{33} &= E[d_3^2] = E[\Delta_2^2 + \Delta_3^2 + 2\Delta_2\Delta_3] \\
&= F_3(0, 2, 0; n - 1) + F_3(0, 0, 2; n - 1) + 2F_3(0, 1, 1; n - 1), \\
a_{44} &= E[d_4^2] = E[\Delta_3^2] = F_3(0, 0, 2; n - 1), \\
a_{12} &= a_{21} = E[d_2] = F_3(1, 0, 0; n - 1) + F_3(0, 1, 0; n - 1) + F_3(0, 0, 1; n - 1), \\
a_{13} &= a_{31} = E[d_3] = F_3(0, 1, 0; n - 1) + F_3(0, 0, 1; n - 1), \\
a_{14} &= a_{41} = E[d_4] = F_3(0, 0, 1; n - 1), \\
a_{23} &= a_{32} = a_{24} = a_{42} = a_{34} = a_{43} = 0.
\end{aligned}$$

References

- [1] Dey, D. K. and Srinivasan, C. (1985). Estimation of a covariance matrix under Stein's loss. *Ann. Statist.*, **13**, 1581–1591.
- [2] James, W. and Stein, C. (1961). Estimation with quadratic loss. *Proc. Fourth Berkeley Symp. Math. Statist. Prob.*, **1**, 361–380. Univ. of California Press.
- [3] Krishnamoorthy, K. and Gupta, A. K. (1989). Improved minimax estimation of a normal precision matrix. *Canad. J. Statist.*, **17**, 91–102.
- [4] Lin, S. P. and Perlman, M. D. (1985). A Monte Carlo comparison of four estimators for a covariance matrix. *Multivariate Analysis*, **6**, 411–429, ed. P. R. Krishnaiah, North Holland, Amsterdam.
- [5] Muirhead, R. J. (1982). *Aspects of Multivariate Statistical Theory*, Wiley, New York.
- [6] Olkin, I. and Selliah, J. B. (1977). Estimating covariances in a multivariate normal distribution. *Statistical Decision Theory and Related Topics*, **2**, 313–326, ed. Gupta, S. S. and Moore, D., Academic Press, New York.

- [7] Pal, N. (1993). Estimating the normal dispersion matrix and the precision matrix from a decision-theoretic point of view: a review. *Statist. Papers*, **34**, 1-26.
- [8] Sheena, Y. (2002). On minimaxity of some orthogonally invariant estimators of bivariate normal dispersion matrix. *J. Japan Statist. Soc.*, **32**, 193–207.
- [9] Sheena, Y. and Takemura, A. (1992). Inadmissibility of non-order-preserving orthogonally invariant estimators of the covariance matrix in the case of Stein’s loss. *Journal of Multivariate Analysis*, **41**, 117–131.
- [10] Sheena, Y. and Takemura, A. (2006). An asymptotic expansion of Wishart distribution when the population eigenvalues are infinitely dispersed. *Statistical Methodology*, doi:10.1016/j.stamet.2006.05.001.
- [11] Stein, C. (1977). Estimating the covariance matrix. Unpublished manuscript.
- [12] Sugiura, N. and Ishibayashi, H. (1997). Reference prior Bayes estimator for bivariate normal covariance matrix with risk comparison. *Comm. Statist.–Theory Meth.*, **26**, 2203-2221.
- [13] Takemura, A. and Sheena, Y. (2005). Distribution of eigenvalues and eigenvectors of Wishart matrix when the population eigenvalues are infinitely dispersed and its application to minimax estimation of covariance matrix. *Journal of Multivariate Analysis*, **94**, 271-299.