

**MATHEMATICAL ENGINEERING  
TECHNICAL REPORTS**

**Application of Arrangement Theory to  
Unfolding Models**

Hidehiko KAMIYA, Akimichi TAKEMURA and  
Norihide TOKUSHIGE

METR 2010-07

March 2010

DEPARTMENT OF MATHEMATICAL INFORMATICS  
GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY  
THE UNIVERSITY OF TOKYO  
BUNKYO-KU, TOKYO 113-8656, JAPAN

**WWW page: <http://www.keisu.t.u-tokyo.ac.jp/research/techrep/index.html>**

The METR technical reports are published as a means to ensure timely dissemination of scholarly and technical work on a non-commercial basis. Copyright and all rights therein are maintained by the authors or by other copyright holders, notwithstanding that they have offered their works here electronically. It is understood that all persons copying this information will adhere to the terms and constraints invoked by each author's copyright. These works may not be reposted without the explicit permission of the copyright holder.

# Application of arrangement theory to unfolding models

Hidehiko Kamiya, Akimichi Takemura and Norihide Tokushige

## Abstract.

Arrangement theory plays an essential role in the study of the unfolding model used in many fields. This paper describes how arrangement theory can be usefully employed in solving the problems of counting (i) the number of admissible rankings in an unfolding model and (ii) the number of ranking patterns generated by unfolding models. The paper is mostly expository but also contains some new results such as simple upper and lower bounds for the number of ranking patterns in the unidimensional case.

## §1. Introduction

The unfolding model (Coombs [6], De Leeuw [8]) is a model for preference rankings in psychometrics. It is now widely applied not only in psychometrics (De Soete, Feger and Klauer [10]) but also in other fields such as marketing science (DeSarbo and Hoffman [9]) and voting theory (Clinton, Jackman and Rivers [5]). The model is also used as a submodel for more complex models, as in item response theory for unfolding (Andrich [1, 2]). Moreover, in the context of Voronoi diagrams, this model can be regarded as a higher-order Voronoi diagram (Okabe, Boots, Sugihara and Chiu [22]).

The unfolding model describes the ranking process in which judges rank a set of objects in order of preference. In this model, judges and objects are assumed to be represented by points in the Euclidean space  $\mathbb{R}^n$ . Suppose a judge  $y \in \mathbb{R}^n$  ranks  $m$  objects  $x_1, \dots, x_m \in \mathbb{R}^n$ . According to the unfolding model,  $y$  ranks  $x_1, \dots, x_m$  in descending order of

---

2000 *Mathematics Subject Classification.* 32S22, 52C35, 62F07.

*Key words and phrases.* all-subset arrangement, braid arrangement, chamber, characteristic polynomial, finite field method, hyperplane arrangement, intersection poset, mid-hyperplane arrangement, partition lattice, ranking pattern, unfolding model.

proximity in the usual Euclidean distance. Hence,  $y$  likes  $x_{i_1}$  best,  $x_{i_2}$  second best, and so on, iff  $\|y - x_{i_1}\| < \|y - x_{i_2}\| < \cdots < \|y - x_{i_m}\|$ . In this case, we will say  $y$  gives ranking  $(i_1 i_2 \cdots i_m)$ .

For a given  $m$ -tuple  $(x_1, \dots, x_m)$  of objects, let  $\text{RP}^{\text{UF}}(x_1, \dots, x_m)$  be the set of admissible rankings, i.e.,  $(i_1 \cdots i_m)$  such that  $\|y - x_{i_1}\| < \cdots < \|y - x_{i_m}\|$  for some  $y \in \mathbb{R}^n$ . We call  $\text{RP}^{\text{UF}}(x_1, \dots, x_m)$  the ranking pattern of the unfolding model with  $m$ -tuple  $(x_1, \dots, x_m)$ . In the psychometric literature, there has not been much study on the structure of the ranking pattern. In this paper, we investigate the ranking pattern by using the theory of hyperplane arrangements (Orlik and Terao [23]). Specifically, we consider the following two problems:

- (i) Find the cardinality of  $\text{RP}^{\text{UF}}(x_1, \dots, x_m)$  for a given generic  $m$ -tuple  $(x_1, \dots, x_m)$ ;
- (ii) Find the cardinality of

$$\{\text{RP}^{\text{UF}}(x_1, \dots, x_m) : (x_1, \dots, x_m) \text{ is a generic } m\text{-tuple}\}.$$

The first problem asks how many rankings are admissible in one unfolding model, and the second inquires how many ranking patterns are possible by using different unfolding models (that is, by taking different choices of  $m$ -tuples of objects). As we will see, these problems can be reduced to those of counting the numbers of chambers of some real arrangements; moreover, the latter problems can be solved by employing general results in the theory of hyperplane arrangements (e.g., Zaslavsky's result on the number of chambers of a real arrangement, the finite field method, etc.). In this sense, arrangement theory plays an essential role in the study of the unfolding model.

This paper gives a survey of recent results ([13], [14], [15], [19]) on the problems stated above. It also contains new results on upper and lower bounds for the number of ranking patterns in the unidimensional case  $n = 1$ . In addition, the problem of counting inequivalent ranking patterns (i.e., those which cannot be obtained from one another by just the relabeling of the objects) when  $n = 1$  was not dealt with specifically in [13] but is discussed fully in the present paper.

The organization of the paper is as follows. In Section 2, we define genericness of the unfolding model, and give the answer to problem (i) above, i.e., the number of admissible rankings of the unfolding model with generic objects. Next, in Section 3 we discuss the problem of counting the number of ranking patterns (problem (ii)). In Subsection 3.1, we deal with the unidimensional case, and give the number of ranking patterns in terms of the number of chambers of the mid-hyperplane arrangement. We also provide explicit upper and lower bounds for the

number of ranking patterns. In Subsection 3.2, we treat the unfolding model of codimension one, where the restriction by dimension is weakest. In this case, we describe how the number of ranking patterns can be expressed by the number of chambers of an arrangement called the all-subset arrangement.

## §2. Number of admissible rankings

In this section, we define genericness of the unfolding model, and discuss the problem of counting the number of admissible rankings generated by the unfolding model with generic objects.

Suppose we are given  $x_1, \dots, x_m \in \mathbb{R}^n$  with  $m \geq 3$  and  $n \leq m - 2$ .

In general, for  $m$  distinct points  $z_1, \dots, z_m \in \mathbb{R}^\nu$  ( $m \geq \nu + 1$ ), let  $\overline{z_i z_j}$  denote the one-simplex connecting two points  $z_i$  and  $z_j$  ( $i < j$ ). Consider the following condition:

- (A) The union of  $\nu$  distinct one-simplices  $\overline{z_{i_k} z_{j_k}}$  ( $i_k < j_k$ ,  $k = 1, \dots, \nu$ ) contains no loop if and only if the corresponding vectors  $z_{i_k} - z_{j_k}$  ( $k = 1, \dots, \nu$ ) are linearly independent.

We assume  $x_1, \dots, x_m \in \mathbb{R}^n$  ( $n \leq m - 2$ ) are generic in the sense that they satisfy the following two conditions:

- (A1) The  $m$  points  $x_1, \dots, x_m \in \mathbb{R}^n$  satisfy condition (A).  
 (A2) The  $m$  points  $(x_1^T, \|x_1\|^2)^T, \dots, (x_m^T, \|x_m\|^2)^T \in \mathbb{R}^{n+1}$  satisfy condition (A).

Now, according to the unfolding model, judge  $y \in \mathbb{R}^n$  prefers  $x_i$  to  $x_j$  ( $i \neq j$ ) iff  $\|y - x_i\| < \|y - x_j\|$ . This condition is equivalent to  $y$  being on the same side as  $x_i$  of the perpendicular bisector

$$\begin{aligned} H_{ij} &:= \{y \in \mathbb{R}^n : \|y - x_i\| = \|y - x_j\|\} \\ &= \{y \in \mathbb{R}^n : (x_i - x_j)^T (y - \frac{x_i + x_j}{2}) = 0\} \end{aligned}$$

of the line segment  $\overline{x_i x_j}$  joining  $x_i$  and  $x_j$ . Let us define a hyperplane arrangement

$$\mathcal{A}_{m,n} = \mathcal{A}_{m,n}(x_1, \dots, x_m) := \{H_{ij} : 1 \leq i < j \leq m\}$$

in  $\mathbb{R}^n$ . We call  $\mathcal{A}_{m,n}$  the unfolding arrangement.

Then  $\mathcal{A}_{m,n}$ , like any real hyperplane arrangement, cuts  $\mathbb{R}^n$  into chambers, i.e., connected components of the complement  $\mathbb{R}^n \setminus \bigcup \mathcal{A}_{m,n}$ , where  $\bigcup \mathcal{A}_{m,n} := \bigcup_{H \in \mathcal{A}_{m,n}} H$ . Moreover, each of these chambers is of the form

$$C_{i_1 \dots i_m} := \{\|y - x_{i_1}\| < \dots < \|y - x_{i_m}\|\} \neq \emptyset$$

for some admissible ranking  $(i_1 \cdots i_m) \in \mathbb{P}_m$ , where  $\mathbb{P}_m$  denotes the set of permutations of  $[m] := \{1, \dots, m\}$ .

We observe that  $y \in \mathbb{R}^n$  gives ranking  $(i_1 \cdots i_m) \in \mathbb{P}_m$  if and only if  $y \in C_{i_1 \cdots i_m} \neq \emptyset$ . Thus there is a one-to-one correspondence between the set of admissible rankings and the set of chambers  $\mathbf{Ch}(\mathcal{A}_{m,n})$  of  $\mathcal{A}_{m,n}$ :

$$(i_1 \cdots i_m) \leftrightarrow C_{i_1 \cdots i_m}$$

for  $(i_1 \cdots i_m)$  such that  $C_{i_1 \cdots i_m} \neq \emptyset$ . This implies that the problem of counting the number of admissible rankings reduces to that of counting the number of chambers of  $\mathcal{A}_{m,n}$ . The answer to the latter problem is given by the theorem below. Let  $\mathcal{S}_k^m$  ( $k \in \mathbb{Z}$ ) be the signless Stirling numbers of the first kind:  $t(t+1) \cdots (t+m-1) = \sum_k \mathcal{S}_k^m t^k$ .

**Theorem 1** (Good and Tideman [11], Kamiya and Takemura [14, 15], Zaslavsky [29]). *Suppose  $x_1, \dots, x_m \in \mathbb{R}^n$  ( $n \leq m-2$ ) are generic. Then, the number of chambers of  $\mathcal{A}_{m,n} = \mathcal{A}_{m,n}(x_1, \dots, x_m)$  is*

$$|\mathbf{Ch}(\mathcal{A}_{m,n})| = \mathcal{S}_{m-n}^m + \mathcal{S}_{m-n+1}^m + \cdots + \mathcal{S}_m^m.$$

Furthermore, the number of bounded chambers of  $\mathcal{A}_{m,n}$  is

$$\mathcal{S}_{m-n}^m - \mathcal{S}_{m-n+1}^m + \mathcal{S}_{m-n+2}^m - \cdots + (-1)^n \mathcal{S}_m^m.$$

The proof of Theorem 1 is based on Zaslavsky's general result on the number of chambers of an arrangement (Zaslavsky [28]) and the following proposition. Denote by  $\Pi_m$  the partition lattice, consisting of partitions of  $[m]$  and ordered by refinement. Further, let  $\Pi_m^n$  stand for the rank  $n$  truncation of  $\Pi_m$ , i.e., the subposet of  $\Pi_m$  comprising elements of rank ( $= m - \#$  of blocks) at most  $n$ .

**Proposition 1** (Kamiya and Takemura [14, 15]). *The intersection poset  $L(\mathcal{A}_{m,n})$  of the unfolding arrangement  $\mathcal{A}_{m,n}$  is isomorphic to  $\Pi_m^n$ :*

$$L(\mathcal{A}_{m,n}) \cong \Pi_m^n.$$

The isomorphism is given by

$$L(\mathcal{A}_{m,n}) \ni X \mapsto I_X \in \Pi_m^n,$$

where  $I_X$  is the partition of  $[m]$  into equivalence classes under the equivalence relation  $\sim_X$  defined by  $i \sim_X j \iff X \subseteq H_{ij}$  ( $H_{ii} := \mathbb{R}^n$ ).

**Remark 1.** *When  $n \geq m-1$ , and  $x_1, \dots, x_m \in \mathbb{R}^n$  satisfy condition (A1) with the  $\nu = n$  in (A) replaced by  $m-1$ , we can easily see that  $|\mathbf{Ch}(\mathcal{A}_{m,n})| = m!$  and that the number of bounded chambers of  $\mathcal{A}_{m,n}$  is zero (so the results in Theorem 1 continue to be valid). Therefore, all  $m!$  rankings arise as unbounded chambers of  $\mathcal{A}_{m,n}$  in this case.*

### §3. Number of ranking patterns

In this section, we consider the problem of counting the number of ranking patterns. We treat two extreme cases—the unidimensional unfolding model:  $n = 1$  (Subsection 3.1) and the unfolding model of codimension one:  $n = m - 2$  (Subsection 3.2).

#### 3.1. Unidimensional unfolding models

In this subsection, we look into the problem of counting the number of ranking patterns of unidimensional unfolding models:  $n = 1$ . A related problem is studied in Stanley [24].

In this case  $n = 1$ , objects are  $m$  points on the real line:  $x_1, \dots, x_m \in \mathbb{R}$ . We assume  $x_1, \dots, x_m$  are generic, i.e., the midpoints  $x_{ij} := (x_i + x_j)/2$ ,  $1 \leq i < j \leq m$ , are all distinct. This condition can be written as

$$(x_1, \dots, x_m) \in \mathbb{R}^m \setminus \bigcup \mathcal{M}_m,$$

where  $\mathcal{M}_m := \mathcal{B}_m \cup \mathcal{N}_m$  is the mid-hyperplane arrangement (Kamiya, Orlik, Takemura and Terao [13]) with

$$\begin{aligned} \mathcal{B}_m &:= \{K_{ij} : 1 \leq i < j \leq m\}, & K_{ij} &:= \{(x_1, \dots, x_m) \in \mathbb{R}^m : x_i = x_j\}, \\ \mathcal{N}_m &:= \{H_{ijkl} : (i, j, k, l) \in I_4\}, \\ H_{ijkl} &:= \{(x_1, \dots, x_m) \in \mathbb{R}^m : x_i + x_j = x_k + x_l\}, \\ I_4 &:= \{(i, j, k, l) : i, j, k, l \text{ are all distinct,} \\ &\quad 1 \leq i < j \leq m, i < k < l \leq m\}. \end{aligned}$$

(In this subsection, we write elements of  $\mathbb{R}^m$  as row vectors.) Note that  $\mathcal{B}_m$  is the braid arrangement. We have  $H_{ij} = \{x_{ij}\}$ ,  $1 \leq i < j \leq m$ , and  $\mathcal{A}_{m,1} = \{\{x_{ij}\} : 1 \leq i < j \leq m\}$ .

An  $m$ -tuple  $\mathbf{x} := (x_1, \dots, x_m)$  of objects gives the ranking pattern

$$\text{RP}^{\text{UF}}(\mathbf{x}) = \{(i_1 \cdots i_m) \in \mathbb{P}_m : |y - x_{i_1}| < \cdots < |y - x_{i_m}| \text{ for some } y \in \mathbb{R}\}.$$

We want to know

$$(1) \quad r(m) := |\{\text{RP}^{\text{UF}}(\mathbf{x}) : \mathbf{x} \in \mathbb{R}^m \setminus \bigcup \mathcal{M}_m\}|.$$

The braid arrangement  $\mathcal{B}_m$  has a chamber  $C_0 \in \mathbf{Ch}(\mathcal{B}_m)$  defined by  $x_1 < \cdots < x_m$ :

$$C_0 := \{(x_1, \dots, x_m) \in \mathbb{R}^m : x_1 < \cdots < x_m\}.$$

Let us concentrate our attention on  $C_0$ . For  $\mathbf{x} = (x_1, \dots, x_m) \in C_0 \setminus \bigcup \mathcal{N}_m$  and  $\mathbf{x}' = (x'_1, \dots, x'_m) \in C_0 \setminus \bigcup \mathcal{N}_m$ , we can easily see that

$\text{RP}^{\text{UF}}(\mathbf{x}) = \text{RP}^{\text{UF}}(\mathbf{x}')$  if and only if the order of the midpoints on  $\mathbb{R}$  is the same for  $\mathbf{x}$  and  $\mathbf{x}'$  (i.e.,  $\forall(i, j, k, l) \in I_4 : x_{ij} < x_{kl} \iff x'_{ij} < x'_{kl}$ ). Noting that  $x_{ij} < x_{kl}$  iff  $(x_1, \dots, x_m) \in H_{ijkl}^- := \{(x_1, \dots, x_m) \in \mathbb{R}^m : x_i + x_j < x_k + x_l\}$ , we obtain the following lemma.

**Lemma 1** (Kamiya, Orlik, Takemura and Terao [13]). *For  $\mathbf{x}, \mathbf{x}' \in C_0 \setminus \bigcup \mathcal{N}_m$ , we have  $\text{RP}^{\text{UF}}(\mathbf{x}) = \text{RP}^{\text{UF}}(\mathbf{x}')$  if and only if  $\mathbf{x}$  and  $\mathbf{x}'$  are in the same chamber of  $\mathcal{N}_m$ .*

Put

$$r_0(m) := |\{\text{RP}^{\text{UF}}(\mathbf{x}) : \mathbf{x} \in C_0 \setminus \bigcup \mathcal{N}_m\}|,$$

i.e., the number of ranking patterns of unidimensional unfolding models with generic  $m$ -tuples such that  $x_1 < \dots < x_m$ . Then, by Lemma 1 we have

$$(2) \quad r_0(m) = \frac{|\mathbf{Ch}(\mathcal{M}_m)|}{m!}$$

(Kamiya, Orlik, Takemura and Terao [13]).

Now consider  $r(m)$  in (1). For  $\mathbf{x} = (x_1, \dots, x_m) \in \mathbb{R}^m \setminus \bigcup \mathcal{M}_m$ , define  $-\mathbf{x} := (-x_1, \dots, -x_m) \in \mathbb{R}^m \setminus \bigcup \mathcal{M}_m$ . Then, clearly we have  $\text{RP}^{\text{UF}}(\mathbf{x}) = \text{RP}^{\text{UF}}(-\mathbf{x})$ . On the other hand, for  $C, C' \in \mathbf{Ch}(\mathcal{M}_m)$  such that  $C' \neq \pm C$  ( $-C := \{-\mathbf{x} : \mathbf{x} \in C\}$ ), we can easily see that  $\text{RP}^{\text{UF}}(\mathbf{x}) \neq \text{RP}^{\text{UF}}(\mathbf{x}')$  for  $\mathbf{x} \in C$  and  $\mathbf{x}' \in C'$ . These two facts, together with Lemma 1, yield the following theorem.

**Theorem 2.** *The number of ranking patterns of unidimensional unfolding models with generic  $m$ -tuples of objects is*

$$r(m) = \frac{m!}{2} r_0(m) = \frac{|\mathbf{Ch}(\mathcal{M}_m)|}{2}, \quad m \geq 3.$$

Let us define equivalence of ranking patterns by saying that two ranking patterns  $\text{RP}^{\text{UF}}(\mathbf{x})$  and  $\text{RP}^{\text{UF}}(\mathbf{x}')$  are equivalent iff

$$(3) \quad \text{RP}^{\text{UF}}(\mathbf{x}) = \sigma \text{RP}^{\text{UF}}(\mathbf{x}') \quad \text{for some } \sigma \in \mathfrak{S}_m,$$

where  $\mathfrak{S}_m$  is the symmetric group on  $m$  letters, consisting of all bijections:  $[m] \rightarrow [m]$ , and  $\sigma \text{RP}^{\text{UF}}(\mathbf{x}') := \{(\sigma(i_1) \dots \sigma(i_m)) : (i_1 \dots i_m) \in \text{RP}^{\text{UF}}(\mathbf{x}')\}$ . We want to find the number of inequivalent ranking patterns.

Let  $r_{\text{IE}}(m)$  be the number of inequivalent ranking patterns of unidimensional unfolding models with generic  $m$ -tuples of objects:

$$r_{\text{IE}}(m) := |\{[\text{RP}^{\text{UF}}(\mathbf{x})] : \mathbf{x} \in \mathbb{R}^m \setminus \bigcup \mathcal{M}_m\}|,$$



where  $[\cdot]$  stands for the equivalence class under the equivalence relation defined by (3). We will see that  $r_{\text{IE}}(m)$  is half of  $r_0(m)$  for  $m \geq 4$ . Suppose we are given  $\mathbf{x} = (x_1, \dots, x_m) \in C_0 \setminus \bigcup \mathcal{N}_m$  with  $m \geq 4$ . Then  $\mathbf{x}' = (x'_1, \dots, x'_m) := (-x_m, \dots, -x_1)$  also lies in  $C_0 \setminus \bigcup \mathcal{N}_m$ :  $\mathbf{x}' \in C_0 \setminus \bigcup \mathcal{N}_m$ . Moreover, since  $m \geq 4$ , four indices  $1, 2, m-1, m$  are all distinct and we have  $x_{1m} < x_{2,m-1}$  iff  $x'_{1m} > x'_{2,m-1}$ . This means  $\text{RP}^{\text{UF}}(\mathbf{x}) \neq \text{RP}^{\text{UF}}(\mathbf{x}')$  by Lemma 1. However,  $[\text{RP}^{\text{UF}}(\mathbf{x})] = [\text{RP}^{\text{UF}}(\mathbf{x}')] = [\text{RP}^{\text{UF}}(-\mathbf{x})]$  since  $\text{RP}^{\text{UF}}(\mathbf{x}) = \text{RP}^{\text{UF}}(-\mathbf{x})$ . Next, it can be seen that any  $\mathbf{x}'' \in C_0 \setminus \bigcup \mathcal{N}_m$  such that  $\text{RP}^{\text{UF}}(\mathbf{x}'') \neq \text{RP}^{\text{UF}}(\mathbf{x})$  and  $[\text{RP}^{\text{UF}}(\mathbf{x}'')] = [\text{RP}^{\text{UF}}(\mathbf{x})]$  satisfies  $\text{RP}^{\text{UF}}(\mathbf{x}'') = \text{RP}^{\text{UF}}(\mathbf{x}')$ . These arguments lead to the following theorem.

**Theorem 3.** *The number of inequivalent ranking patterns of uni-dimensional unfolding models with generic  $m$ -tuples of objects is*

$$r_{\text{IE}}(m) = \begin{cases} r_0(3) = \frac{|\text{Ch}(\mathcal{B}_3)|}{3!} = 1 & \text{if } m = 3, \\ \frac{r_0(m)}{2} = \frac{|\text{Ch}(\mathcal{M}_m)|}{2 \cdot m!} & \text{if } m \geq 4. \end{cases}$$

So far, we have expressed the number of ranking patterns in terms of the number of chambers of an arrangement. We can use the finite field method (Athanasiadis [3, 4], Crapo and Rota [7], Kamiya, Takemura and Terao [16, 17, 18], Stanley [25, Lecture 5]) to calculate specific values of  $r_0(m)$ ,  $m \leq 10$ :

$$\begin{aligned} r_0(4) &= 2, \quad r_0(5) = 12, \quad r_0(6) = 168, \quad r_0(7) = 4680, \\ r_0(8) &= 229386, \quad r_0(9) = 18330206, \quad r_0(10) = 1120831141. \end{aligned}$$

The values of  $r(m)$  for  $m \leq 8$  are given in Kamiya, Orlik, Takemura and Terao [13] along with the characteristic polynomials  $\chi(\mathcal{M}_m, t)$  of  $\mathcal{M}_m$ ,  $m \leq 8$ . After [13], the second author of the present paper, Takemura [26], calculated  $\chi(\mathcal{M}_9, t)$  and  $r_0(9)$ ; later Ishiwata [12] obtained  $\chi(\mathcal{M}_{10}, t)$  and  $r_0(10)$  after an extensive computation. The characteristic polynomials found by them are:

$$\begin{aligned} \chi(\mathcal{M}_9, t) &= t(t-1)(t^7 - 413t^6 + 73780t^5 - 7387310t^4 + 447514669t^3 \\ &\quad - 16393719797t^2 + 336081719070t - 2972902161600), \\ \chi(\mathcal{M}_{10}, t) &= t(t-1)(t^8 - 674t^7 + 201481t^6 - 34896134t^5 + 3830348179t^4 \\ &\quad - 272839984046t^3 + 12315189583899t^2 \\ &\quad - 321989533359786t + 3732690616086600). \end{aligned}$$

However, for large values of  $m$ , the finite field method is not feasible. We will provide simple upper and lower bounds for  $r_0(m)$ .

**Theorem 4.** *For all  $m \geq 4$ , we have*

$$2 \left( \frac{3}{4} \right)^{m-4} \{(m-3)!\}^2 \leq r_0(m) < \frac{2}{m!} \left\{ \frac{em(m-1)^2}{8} \right\}^{m-2}.$$

*Proof.* First, we derive the upper bound in the theorem.

Define  $H_0 := \{(x_1, \dots, x_m) \in \mathbb{R}^m : x_1 + \dots + x_m = 0\}$ , and consider the essentialization (Stanley [25, p.392])  $\mathcal{M}_m^0 := \{H \cap H_0 : H \in \mathcal{M}_m\}$  of  $\mathcal{M}_m$ . Since  $L(\mathcal{M}_m^0) \cong L(\mathcal{M}_m)$ , we may consider the essential, central arrangement  $\mathcal{M}_m^0$  in  $H_0$  ( $\dim H_0 = m-1$ ) instead of  $\mathcal{M}_m$ .

Recall, in general, that  $h$  hyperplanes divide  $\mathbb{R}^d$  into at most  $\sum_{i=0}^d \binom{h}{i} \leq (eh/d)^d := c(h, d)$  chambers (see, e.g., [20, Proposition 6.1.1] and [21, Theorem 2.6.1]). Thus,  $\tilde{h}$  linear hyperplanes divide  $\mathbb{R}^{\tilde{d}}$  into at most  $2c(\tilde{h}-1, \tilde{d}-1)$  chambers.

In our case,  $\mathcal{M}_m^0$  is central, so we can take  $\tilde{h} = |\mathcal{M}_m| = |\mathcal{B}_m| + |\mathcal{N}_m| = \binom{m}{2} + 3\binom{m}{4} \leq m(m-1)^2(m-2)/8$  ( $m \geq 4$ ) and  $\tilde{d} = m-1$ . Hence, we have

$$\begin{aligned} |\mathbf{Ch}(\mathcal{M}_m^0)| &\leq 2c(\tilde{h}-1, \tilde{d}-1) \\ &\leq 2 \times \left\{ \frac{e \left( \frac{m(m-1)^2(m-2)}{8} - 1 \right)}{m-2} \right\}^{m-2} \\ &< 2 \times \left\{ \frac{em(m-1)^2}{8} \right\}^{m-2}. \end{aligned}$$

This together with (2) and  $|\mathbf{Ch}(\mathcal{M}_m)| = |\mathbf{Ch}(\mathcal{M}_m^0)|$  gives the upper bound of  $r_0(m)$  in the theorem.

Next, we will obtain the lower bound in the theorem.

Let  $\mathbf{x} = (x_1, \dots, x_m)$ ,  $x_1 < \dots < x_m$  be fixed. We add one more object  $y = x_m + 2t$  ( $t > 0$ ) to  $\mathbf{x}$ , and we will count the number of ranking patterns arising from  $\mathbf{y}_t = (\mathbf{x}, y)$ ,  $t > 0$ . Let  $M = \{x_{ij} : 1 \leq i < j \leq m\}$  be the set of midpoints for  $\mathbf{x}$ , and  $Y_t = \{x_{im} + t : 1 \leq i \leq m\}$  the set of midpoints of  $x_i$  ( $1 \leq i \leq m$ ) and  $y$ . Then  $M \cup Y_t$  is the set of midpoints for  $\mathbf{y}_t$ . To guarantee all these midpoints are distinct, we require the following. First, by perturbing each  $x_i$  without changing the ranking pattern of  $\mathbf{x}$ , we may assume that  $x_1, \dots, x_m$  are independent over  $\mathbb{Q}$ . Then we have  $|M \cap Y_t| \leq 1$  for all  $t > 0$ . Next, let  $T_0 = \{t > 0 : |M \cap Y_t| = 1\}$ ,  $T_1 = (0, \infty) \setminus T_0$ , and we only consider  $t \in T_1$ . Then  $M \cup Y_t$  is legal, i.e., all midpoints are distinct.

Now the crucial observation is as follows:  $|\{\mathbf{RP}^{\text{UF}}(\mathbf{y}_t) : t \in T_1\}| = 1 + |T_0|$ . Moreover, we have  $|T_0| = \sum_{i=1}^{m-1} |V_i|$ , where  $V_i = \{v \in M :$

$x_{im} < v$ . Using  $|V_i| \geq m - 1 - i$  obtained by  $V_i \supset \{x_{jm} : i < j < m\}$ , we have

$$|\{\text{RP}^{\text{UF}}(\mathbf{y}_t) : t \in T_1\}| = 1 + \sum_{i=1}^{m-1} |V_i| \geq 1 + |V_1| + \frac{(m-3)(m-2)}{2} =: N.$$

Namely,  $N$  is a lower bound for the number of ranking patterns arising from  $\mathbf{y}_t$ ,  $t \in T_1$ .

Applying exactly the same argument to  $\mathbf{x}' = (-x_m, \dots, -x_1)$  instead of  $\mathbf{x}$ , we see that the number of ranking patterns arising from  $(\mathbf{x}', -x_1 + 2t)$ ,  $t > 0$  (or equivalently,  $(x_1 - 2t, \mathbf{x})$ ,  $t > 0$ ) is at least  $N' = 1 + |V'_1| + (m-3)(m-2)/2$ , where  $|V'_1| = |\{u \in M : u < x_{1m}\}| = \binom{m}{2} - |V_1| - 1$ . Notice that  $N + N' = 1 + \binom{m}{2} + (m-3)(m-2) > (3/2)(m-2)^2$ . Therefore, by the averaging argument, we have

$$r_0(m+1) \geq r_0(m) \times \frac{1}{2}(N + N') > \frac{3}{4}(m-2)^2 r_0(m).$$

So the induction starting from  $r_0(4) = 2$  gives the desired lower bound. Q.E.D.

Let  $\ell(m)$  and  $u(m)$  be the lower and upper bounds in the theorem, respectively. A computation shows  $\{u(m)\}^{1/m}/m^2 \rightarrow e^2/8 \approx 0.92$  and  $\{\ell(m)\}^{1/m}/m^2 \rightarrow 3/(4e^2) \approx 0.1$  as  $m \rightarrow \infty$ . It would be interesting to prove (or disprove) the existence of  $\lim\{r_0(m)\}^{1/m}/m^2$ .

Strangely enough,  $r_0(m) = a(m)$  holds for  $4 \leq m \leq 7$ , where

$$a(m) := \frac{(m-2)\{(m-2)^{m-3} - 1\} \cdot (m-4)!}{m-3},$$

but  $r_0(8) > a(8)$ ,  $r_0(9) > a(9)$ ,  $r_0(10) < a(10)$ . Also,  $a(m)$  satisfies  $\{a(m)\}^{1/m}/m^2 \rightarrow 1/e \approx 0.37$ .

Thrall [27] gave an upper bound  $f(m)$  for  $r_0(m)$ :

$$f(m) := \frac{\{\frac{m(m-1)}{2}\}! \prod_{i=1}^{m-2} i!}{\prod_{i=1}^{m-1} (2i-1)!}.$$

We can see our  $u(m)$  satisfies  $f(m) < u(m)$  for  $m \leq 8$ ,  $f(m) > u(m)$  for  $m \geq 9$ , and  $u(m) = o(f(m))$ .

We list the values of  $r_0(m)$ ,  $a(m)$ ,  $f(m)$  and approximate values of  $\ell(m)$ ,  $u(m)$  for  $m = 4, \dots, 10$  in Table 1. (For  $\ell(m)$ ,  $m \leq 9$ , and  $u(m)$ ,  $m \leq 6$ , we exhibit  $\lceil \ell(m) \rceil$  and  $\lfloor u(m) \rfloor$ , respectively. For  $\ell(10)$ , we display  $\lceil \ell(m) \times 10^{-4} \rceil \times 10^4$ , and similarly using  $\lfloor \cdot \rfloor$  for  $u(m)$ ,  $m \geq 7$ .)

Table 1.  $r_0(m)$ ,  $a(m)$ ,  $\ell(m)$ ,  $u(m)$ ,  $f(m)$ ,  $4 \leq m \leq 10$ .

$m$	$r_0(m)$	$a(m)$	$\ell(m)$	$u(m)$	$f(m)$
4	2	2	2	12	2
5	12	12	6	334	12
6	168	168	41	18,744	286
7	4,680	4,680	486	$1.82 \times 10^6$	33,592
8	229,386	223,920	9,113	$2.76 \times 10^8$	23,178,480
9	18,330,206	16,470,720	246,038	$6.06 \times 10^{10}$	108,995,910,720
10	1,120,831,141	1,725,655,680	$9.05 \times 10^6$	$1.81 \times 10^{13}$	3,973,186,258,569,120

### 3.2. Unfolding models of codimension one

In this subsection, we deal with the problem of counting the number of ranking patterns of unfolding models of codimension one:  $n = m - 2$  (i.e., when the restriction by dimension is weakest).

First, let us forget the unfolding model for a while and consider the ranking patterns of braid slices.

We begin by defining the ranking pattern of a braid slice. For

$$H_0 = \{x = (x_1, \dots, x_m)^T \in \mathbb{R}^m : x_1 + \dots + x_m = 0\},$$

consider the essential arrangement

$$\mathcal{B}_m^0 := \{H \cap H_0 : H \in \mathcal{B}_m\}$$

in  $H_0$ , and write its chambers as

$$B_{i_1 \dots i_m} := \{x = (x_1, \dots, x_m)^T \in H_0 : x_{i_1} > \dots > x_{i_m}\} \in \mathbf{Ch}(\mathcal{B}_m^0)$$

for  $(i_1 \dots i_m) \in \mathbb{P}_m$ . Moreover, define a hyperplane

$$K_v := \{x \in H_0 : v^T x = 1\}$$

in  $H_0$  for each  $v \in \mathbb{S}^{m-2} := \{x \in H_0 : \|x\| = 1\}$ . Now we call the subset

$$\mathbf{RP}(v) := \{(i_1 \dots i_m) \in \mathbb{P}_m : K_v \cap B_{i_1 \dots i_m} \neq \emptyset\}, \quad v \in \mathbb{S}^{m-2},$$

of  $\mathbb{P}_m$  the ranking pattern of the braid slice by  $K_v$ .

Next, let us define genericness of the braid slice as follows. For the all-subset arrangement (Kamiya, Takemura and Terao [19])

$$\mathcal{A}_m := \{H_I : I \subseteq [m], |I| \geq 1\}$$

with  $H_I := \{x = (x_1, \dots, x_m)^T \in \mathbb{R}^m : \sum_{i \in I} x_i = 0\}$ ,  $\emptyset \neq I \subseteq [m]$ , consider its restriction to  $H_0 = H_{[m]}$ :

$$\begin{aligned} \mathcal{A}_m^0 &:= \mathcal{A}_m^{H_0} = \{H_I^0 : I \subset [m], 1 \leq |I| \leq m-1\}, \\ H_I^0 &:= H_I \cap H_0 \quad (1 \leq |I| \leq m-1). \end{aligned}$$

Then define

$$\mathcal{V} := (H_0 \setminus \bigcup \mathcal{A}_m^0) \cap \mathbb{S}^{m-2}.$$

We will say  $v \in \mathbb{S}^{m-2}$ , or the braid slice by  $K_v$ , is generic if  $v \in \mathcal{V}$ .

Now, we will see that the set of ranking patterns  $\text{RP}(v)$  for generic  $v$ 's is in one-to-one correspondence with the set of chambers of  $\mathcal{A}_m^0$ . Write  $\mathcal{V}$  as  $\mathcal{V} = \bigsqcup_{D \in \mathbf{D}(\mathcal{A}_m^0)} D$  (disjoint union), where

$$\mathbf{D}(\mathcal{A}_m^0) := \{D = \tilde{D} \cap \mathbb{S}^{m-2} : \tilde{D} \in \mathbf{Ch}(\mathcal{A}_m^0)\},$$

which clearly is in one-to-one correspondence with  $\mathbf{Ch}(\mathcal{A}_m^0)$ . Then, we can prove (Kamiya, Takemura and Terao [19]) that there is a bijection from  $\mathbf{D}(\mathcal{A}_m^0)$  to  $\{\text{RP}(v) : v \in \mathcal{V}\}$  given by

$$(4) \quad \mathbf{D}(\mathcal{A}_m^0) \ni D \mapsto \text{RP}(v), \quad v \in D.$$

Hence,

$$\text{RP}_D := \text{RP}(v) \text{ for } v \in D \in \mathbf{D}(\mathcal{A}_m^0)$$

is well-defined, and the mapping  $\mathbf{D}(\mathcal{A}_m^0) \rightarrow \{\text{RP}_D : D \in \mathbf{D}(\mathcal{A}_m^0)\} = \{\text{RP}(v) : v \in \mathcal{V}\} : D \mapsto \text{RP}_D$  is bijective.

Let us get back to the unfolding model and consider the ranking pattern of the unfolding model of codimension one.

Suppose we are given  $x_1, \dots, x_m \in \mathbb{R}^n$  with  $n = m - 2 \geq 1$ . We assume  $x_1, \dots, x_m$  are generic in the sense that they satisfy (A1) and (A2) in Section 2. We call the unfolding model with such  $x_1, \dots, x_m \in \mathbb{R}^{m-2}$  the unfolding model of codimension one (for the reason stated below). In addition, we will assume without loss of generality that  $x_1, \dots, x_m$  are taken so that  $\sum_{i=1}^m x_i = 0$ ,  $\sum_{i=1}^m \|x_i\|^2/m = 1$ .

We will see that the ranking pattern of the unfolding model of codimension one with  $m$ -tuple  $(x_1, \dots, x_m)$ :

$$(5) \quad \text{RP}^{\text{UF}}(x_1, \dots, x_m) = \{(i_1 \cdots i_m) \in \mathbb{P}_m : \|y - x_{i_1}\| < \cdots < \|y - x_{i_m}\| \text{ for some } y \in \mathbb{R}^{m-2}\}$$

can be expressed as the ranking pattern of a braid slice.

Define

$$W = W(x_1, \dots, x_m) = (w_1, \dots, w_{m-2}) := \begin{pmatrix} x_1^T \\ \vdots \\ x_m^T \end{pmatrix} \in \text{Mat}_{m \times (m-2)}(\mathbb{R}),$$

$$u = u(x_1, \dots, x_m) := -\frac{1}{2} \begin{pmatrix} \|x_1\|^2 - 1 \\ \vdots \\ \|x_m\|^2 - 1 \end{pmatrix} \in \mathbb{R}^m,$$

where  $\text{Mat}_{m \times (m-2)}(\mathbb{R})$  denotes the set of  $m \times (m-2)$  matrices with real entries. For the affine map  $\kappa : \mathbb{R}^{m-2} \rightarrow \mathbb{R}^m$  defined by  $\kappa(y) := Wy + u$ ,  $y \in \mathbb{R}^{m-2}$ , consider the image  $K := \text{im } \kappa = \{\kappa(y) : y \in \mathbb{R}^{m-2}\}$  of  $\kappa$ . Then we have

$$K = u + \text{col } W \subset H_0,$$

where  $\text{col } W$  stands for the column space of  $W$ . Using this  $K$ , we can easily see that  $\text{RP}^{\text{UF}}(x_1, \dots, x_m)$  in (5) can be expressed as

$$(6) \quad \text{RP}^{\text{UF}}(x_1, \dots, x_m) = \{(i_1 \cdots i_m) \in \mathbb{P}_m : K \cap B_{i_1 \cdots i_m} \neq \emptyset\}.$$

We have  $\dim K = \dim H_0 - 1$  and  $u \notin \text{col } W$  by (A1) and (A2), respectively. That is,  $K$  is an affine hyperplane of  $H_0$ . For this reason, we called the unfolding model with generic  $x_1, \dots, x_m \in \mathbb{R}^{m-2}$  the unfolding model of codimension one.

Write the affine hyperplane  $K \subset H_0$  as

$$K = K_{\tilde{v}} := \{x \in H_0 : \tilde{v}^T x = \|\tilde{v}\|^2\}$$

using the orthogonal projection of  $u \in H_0$  on  $(\text{col } W)^\perp := \{x \in H_0 : x^T W = 0\}$ :

$$\tilde{v} := \tilde{v}(x_1, \dots, x_m) = u - \text{proj}_{\text{col } W}(u), \quad u = u(x_1, \dots, x_m),$$

where  $\text{proj}_{\text{col } W}$  denotes the orthogonal projection on  $\text{col } W$ . Noting  $\tilde{v} \neq 0$ , we can represent (6) as

$$(7) \quad \text{RP}^{\text{UF}}(x_1, \dots, x_m) = \{(i_1 \cdots i_m) \in \mathbb{P}_m : K_{\mathbf{v}(x_1, \dots, x_m)} \cap B_{i_1 \cdots i_m} \neq \emptyset\},$$

$$\mathbf{v}(x_1, \dots, x_m) := \frac{1}{\|\tilde{v}\|} \tilde{v} \in \mathbb{S}^{m-2},$$

in terms of  $K_{\mathbf{v}(x_1, \dots, x_m)} = \{x \in H_0 : \mathbf{v}(x_1, \dots, x_m)^T x = 1\}$  instead of  $K = K_{\tilde{v}}$ . The right-hand side of (7) is the ranking pattern of the braid slice by  $K_{\mathbf{v}(x_1, \dots, x_m)}$ :  $\text{RP}(\mathbf{v}(x_1, \dots, x_m))$ . Besides, it can be seen that  $\mathbf{v}(x_1, \dots, x_m) \in \mathcal{V}$ .

**Proposition 2** (Kamiya, Takemura and Terao [19]). *For generic  $x_1, \dots, x_m \in \mathbb{R}^{m-2}$ , we have  $v(x_1, \dots, x_m) \in \mathcal{V}$  and*

$$\mathrm{RP}^{\mathrm{UF}}(x_1, \dots, x_m) = \mathrm{RP}(v(x_1, \dots, x_m)).$$

Proposition 2 and bijection (4) tell us that in order to find the number of ranking patterns of unfolding models of codimension one, we need to study the image of the mapping  $v : \{(x_1, \dots, x_m) : x_1, \dots, x_m \in \mathbb{R}^{m-2} \text{ are generic}\} \rightarrow \mathcal{V} = \bigsqcup_{D \in \mathbf{D}(\mathcal{A}_m^0)} D, (x_1, \dots, x_m) \mapsto v(x_1, \dots, x_m)$ . In their main theorem (Theorem 4.1), Kamiya, Takemura and Terao [19] proved that the image  $\mathrm{im} v$  is given by

$$(8) \quad \mathrm{im} v = \mathcal{V}_2 \sqcup D_1 \sqcup \dots \sqcup D_m = \mathcal{V} \setminus ((-D_1) \sqcup \dots \sqcup (-D_m)),$$

where

$$\mathcal{V}_2 := \{v = (v_1, \dots, v_m)^T \in \mathcal{V} : v_j > 0 \text{ for at least two } j \in [m] \text{ and } v_k < 0 \text{ for at least two } k \in [m]\}$$

and

$$\begin{aligned} D_i &:= \{v = (v_1, \dots, v_m)^T \in \mathcal{V} : v_i > 0, v_j < 0 (j \neq i)\} \in \mathbf{D}(\mathcal{A}_m^0), \\ -D_i &:= \{-v : v \in D_i\} \\ &= \{v = (v_1, \dots, v_m)^T \in \mathcal{V} : v_i < 0, v_j > 0 (j \neq i)\} \in \mathbf{D}(\mathcal{A}_m^0) \end{aligned}$$

for  $i \in [m]$ .

By Proposition 2 and  $\mathrm{im} v$  in (8), we obtain the number of ranking patterns of unfolding models of codimension one, which is denoted by

$$q(m) := |\{\mathrm{RP}^{\mathrm{UF}}(x_1, \dots, x_m) : \text{generic } x_1, \dots, x_m \in \mathbb{R}^{m-2}\}|.$$

**Theorem 5** (Kamiya, Takemura and Terao [19]). *The number  $q(m)$  of ranking patterns of unfolding models of codimension one is given by*

$$q(m) = |\mathbf{Ch}(\mathcal{A}_m^0)| - m.$$

Kamiya, Takemura and Terao [19, Lemma 5.3] obtained the characteristic polynomials  $\chi(\mathcal{A}_m^0, t)$  of  $\mathcal{A}_m^0$  for  $m \leq 8$  by the finite field method. Then  $q(m)$  can be calculated by  $q(m) = (-1)^{m-1} \chi(\mathcal{A}_m^0, -1) - m$ :

$$\begin{aligned} q(3) &= 3, \quad q(4) = 28, \quad q(5) = 365, \\ q(6) &= 11286, \quad q(7) = 1066037, \quad q(8) = 347326344 \end{aligned}$$

([19, Corollary 5.5]).

We end this subsection by looking at the problem of finding the number of inequivalent ranking patterns of unfolding models of codimension one.

In (3), we defined equivalence of ranking patterns of unidimensional unfolding models. We define equivalence of ranking patterns of unfolding models of codimension one in an obvious similar manner. At the moment, we can only give an upper bound for the number  $q_{\text{IE}}(m)$  of inequivalent ranking patterns of unfolding models of codimension one:

$$(9) \quad q_{\text{IE}}(m) \leq \frac{|\mathbf{Ch}(\mathcal{A}_m^0 \cup \mathcal{B}_m^0)|}{m!} - 1 = |\mathbf{D}^{1 \cdots m}(\mathcal{A}_m^0)| - 1 = |\mathbf{D}_2^{1 \cdots m}(\mathcal{A}_m^0)| + 1$$

for  $m \geq 3$  (Kamiya, Takemura and Terao [19]), where  $\mathbf{D}^{1 \cdots m}(\mathcal{A}_m^0) := \{D \in \mathbf{D}(\mathcal{A}_m^0) : D \cap B_{1 \cdots m} \neq \emptyset\}$  and  $\mathbf{D}_2^{1 \cdots m}(\mathcal{A}_m^0) := \{D \in \mathbf{D}(\mathcal{A}_m^0) : D \subset \mathcal{V}_2, D \cap B_{1 \cdots m} \neq \emptyset\} = \mathbf{D}^{1 \cdots m}(\mathcal{A}_m^0) \setminus \{D_1, -D_m\}$ . It is shown in [19], however, that the upper bound in (9) is actually the exact number for  $m \leq 6$ . The specific values are

$$q_{\text{IE}}(3) = 1, \quad q_{\text{IE}}(4) = 3, \quad q_{\text{IE}}(5) = 11, \quad q_{\text{IE}}(6) = 55$$

([19, Subsection 6.2]).

**Open problem:** Does the upper bound in (9) agree with the exact number  $q_{\text{IE}}(m)$  for all  $m$ ?

**Acknowledgments.** The first author is supported by JSPS KAKENHI (19540131). The second author is supported by JST CREST. The last author is supported by JSPS KAKENHI (20340022).

## References

- [1] D. Andrich, The application of an unfolding model of the PIRT type to the measurement of attitude, *Appl. Psychol. Meas.*, **12** (1988), 33–51.
- [2] D. Andrich, A probabilistic IRT model for unfolding preference data, *Appl. Psychol. Meas.*, **13** (1989), 193–216.
- [3] C. A. Athanasiadis, Algebraic Combinatorics of Graph Spectra, Subspace Arrangements and Tutte Polynomials, Ph.D. thesis, MIT, 1996.
- [4] C. A. Athanasiadis, Characteristic polynomials of subspace arrangements and finite fields, *Adv. Math.*, **122** (1996), 193–233.
- [5] J. Clinton, S. Jackman and D. Rivers, The statistical analysis of roll call data, *Amer. Polit. Sci. Rev.*, **98** (2004), 355–370.
- [6] C. H. Coombs, Psychological scaling without a unit of measurement, *Psychol. Rev.*, **57** (1950), 145–158.



- [7] H. Crapo and G.-C. Rota, *On the Foundations of Combinatorial Theory: Combinatorial Geometries*, preliminary edition, MIT Press, Cambridge, MA, 1970.
- [8] J. De Leeuw, Multidimensional unfolding, In: *Encyclopedia of Statistics in Behavioral Science*, (eds. in chf. B.S. Everitt, D.C. Howell), John Wiley & Sons, Chichester, West Sussex, 2005, pp. 1289–1294.
- [9] W. S. DeSarbo and D. L. Hoffman, Constructing MDS joint spaces from binary choice data: A multidimensional unfolding threshold model for marketing research, *J. Marketing Res.*, **24** (1987), 40–54.
- [10] G. De Soete, H. Feger and K. C. Klauer, *New Developments in Psychological Choice Modeling*, North-Holland, Amsterdam, 1989.
- [11] I. J. Good and T. N. Tideman, Stirling numbers and a geometric structure from voting theory, *J. Combin. Theory A*, **23** (1977), 34–45.
- [12] M. Ishiwata, *Mid-Hyperplane Arrangement and Ranking Patterns*, (in Japanese), Master’s thesis, Tokyo Metropolitan University, 2007.
- [13] H. Kamiya, P. Orlik, A. Takemura and H. Terao, Arrangements and ranking patterns, *Ann. Combin.*, **10** (2006), 219–235.
- [14] H. Kamiya and A. Takemura, On rankings generated by pairwise linear discriminant analysis of  $m$  populations, *J. Multivariate Anal.*, **61** (1997), 1–28.
- [15] H. Kamiya and A. Takemura, Characterization of rankings generated by linear discriminant analysis, *J. Multivariate Anal.*, **92** (2005), 343–358.
- [16] H. Kamiya, A. Takemura and H. Terao, Periodicity of hyperplane arrangements with integral coefficients modulo positive integers, *J. Algebraic Combin.*, **27** (2008), 317–330.
- [17] H. Kamiya, A. Takemura and H. Terao, The characteristic quasi-polynomials of the arrangements of root systems and mid-hyperplane arrangements, In: *Arrangements, Local Systems and Singularities*, CIMPA Summer School, Galatasaray University, Istanbul, 2007, (eds. F. El Zein, A. I. Suci, M. Tosun, A. M. Uludağ and S. Yuzvinsky), *Progress in Mathematics*, **283**, Birkhäuser Verlag, Basel, 2009, pp. 177–190.
- [18] H. Kamiya, A. Takemura and H. Terao, Periodicity of non-central integral arrangements modulo positive integers, arXiv:0803.2755v1 [math.CO], to appear in *Ann. Combin.*
- [19] H. Kamiya, A. Takemura and H. Terao, Ranking patterns of unfolding models of codimension one, arXiv:1003.0040v1 [math.CO].
- [20] J. Matoušek, *Lectures on Discrete Geometry*, Graduate Texts in Mathematics, **212**, Springer-Verlag, New York, 2002.
- [21] J. Matoušek and J. Nešetřil, *Invitation to Discrete Mathematics*, The Clarendon Press, Oxford University Press, New York, 1998.
- [22] A. Okabe, B. Boots, K. Sugihara and S.N. Chiu, *Spatial Tessellations: Concepts and Applications of Voronoi Diagrams*, 2nd ed., John Wiley & Sons, Chichester, West Sussex, 2000.
- [23] P. Orlik and H. Terao, *Arrangements of Hyperplanes*, Springer-Verlag, Berlin, 1992.

- [24] R. Stanley, Ordering events in Minkowski space, *Adv. Appl. Math.*, **37** (2006), 514–525.
- [25] R. Stanley, An introduction to hyperplane arrangements, In: *Geometric Combinatorics*, (eds. E. Miller, V. Reiner and B. Sturmfels), IAS/Park City Mathematics Series, **13**, American Mathematical Society, Providence, RI, 2007, pp. 389–496.
- [26] A. Takemura, unpublished results, 2005.
- [27] R. M. Thrall, A combinatorial problem, *Michigan Math. J.*, **1** (1952), 81–88.
- [28] T. Zaslavsky, Facing up to Arrangements: Face-Count Formulas for Partitions of Space by Hyperplanes, *Mem. Amer. Math. Soc.*, **154**, 1975.
- [29] T. Zaslavsky, Perpendicular dissection of space, *Discrete Comput. Geom.*, **27** (2002), 303–351.

Hidehiko Kamiya  
*Faculty of Economics*  
*Okayama University*  
*Okayama 700-8530*  
*Japan*  
*E-mail address: kamiya@e.okayama-u.ac.jp*

Akimichi Takemura  
*Graduate School of Information Science and Technology*  
*University of Tokyo*  
*Tokyo 113-0033*  
*Japan*  
*E-mail address: takemura@stat.t.u-tokyo.ac.jp*

Norihide Tokushige  
*College of Education*  
*Ryukyu University*  
*Nishihara, Okinawa 903-0213*  
*Japan*  
*E-mail address: hide@edu.u-ryukyu.ac.jp*