

**MATHEMATICAL ENGINEERING  
TECHNICAL REPORTS**

**Exact Bounds for Steepest Descent  
Algorithms of L-convex Function Minimization**

Kazuo MUROTA and Akiyoshi SHIOURA

METR 2013-24

September 2013

DEPARTMENT OF MATHEMATICAL INFORMATICS  
GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY  
THE UNIVERSITY OF TOKYO  
BUNKYO-KU, TOKYO 113-8656, JAPAN

**WWW page: <http://www.keisu.t.u-tokyo.ac.jp/research/techrep/index.html>**

The METR technical reports are published as a means to ensure timely dissemination of scholarly and technical work on a non-commercial basis. Copyright and all rights therein are maintained by the authors or by other copyright holders, notwithstanding that they have offered their works here electronically. It is understood that all persons copying this information will adhere to the terms and constraints invoked by each author's copyright. These works may not be reposted without the explicit permission of the copyright holder.

# Exact Bounds for Steepest Descent Algorithms of L-convex Function Minimization\*

Kazuo MUROTA

Graduate School of Information Science and Technology  
University of Tokyo, Tokyo 113-8656, Japan

and

Akiyoshi SHIOURA

Graduate School of Information Sciences  
Tohoku University, Sendai 980-8579, Japan

September 2013

## Abstract

We analyze minimization algorithms for  $L^h$ -convex functions in discrete convex analysis, and establish exact bounds for the number of iterations required by the steepest descent algorithm and its variants.

---

\*This research is partially supported by KAKENHI (21360045, 21740060, 24500002) and the Aihara Project, the FIRST program from JSPS.

# 1 Results

In this paper, we discuss minimization algorithms for discrete convex functions defined on integer lattice points called  $L^{\natural}$ -convex functions. Function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  is said to be  $L^{\natural}$ -convex [14] if for every  $p, q \in \text{dom } g$  and every nonnegative  $\lambda \in \mathbb{Z}_+$ , it holds that

$$g(p) + g(q) \geq g((p + \lambda \mathbf{1}) \wedge q) + g(p \vee (q - \lambda \mathbf{1})), \quad (1)$$

where  $\text{dom } g = \{p \in \mathbb{Z}^n \mid g(p) < +\infty\}$ ,  $\mathbf{1} = (1, 1, \dots, 1)$ , and for  $p, q \in \mathbb{Z}^n$  the vectors  $p \wedge q$  and  $p \vee q$  denote, respectively, the vectors of component-wise minimum and maximum of  $p$  and  $q$ . The concept of  $L^{\natural}$ -convex function plays a primary role in the theory of discrete convex analysis [14], and there exist many examples of  $L^{\natural}$ -convex functions arising from applications in various research areas such as discrete optimization, iterative auctions, and computer vision (see Section 2; see also [14]). Some applications of  $L^{\natural}$ -convex functions in inventory systems can also be found in [11, 19].

We consider minimization of an  $L^{\natural}$ -convex function  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  with  $\text{argmin } g \neq \emptyset$ . It is known that this problem can be solved by the following steepest descent algorithm [15], where an initial vector  $p^\circ \in \text{dom } g$  is assumed to be given. Let  $N = \{1, 2, \dots, n\}$  and denote by  $\chi_X \in \{0, 1\}^n$  the characteristic vector of  $X \subseteq N$ , i.e.,  $\chi_X(i) = 1$  if  $i \in X$  and  $\chi_X(i) = 0$  if  $i \in N \setminus X$ .

## Algorithm STEEPESTDESCENT

Step 0: Set  $p := p^\circ$ .

Step 1: Find  $\sigma \in \{+1, -1\}$  and  $X \subseteq N$  that minimize  $g(p + \sigma \chi_X)$ .

Step 2: If  $g(p + \sigma \chi_X) = g(p)$ , then output  $p$  and stop.

Step 3: Set  $p := p + \sigma \chi_X$  and go to Step 1.

**Theorem 1.1** ([15]). *Suppose that  $\text{dom } g$  is bounded. Then, the algorithm STEEPESTDESCENT outputs a minimizer of  $g$  in  $O(nK_\infty)$  iterations, where  $K_\infty = \max\{\|p - q\|_\infty \mid p, q \in \text{dom } g\}$ .*

The bound  $O(nK_\infty)$  for the number of iterations is later improved to  $2K_\infty + 1$  [13, Th. 2.8].

The main aim of this paper is to give a refined analysis of this algorithm in terms of the “distance” between the initial vector and a minimizer of  $g$ . For a vector  $q \in \mathbb{Z}^n$ , denote

$$\|q\|_\infty^+ = \max_{i \in N} \max(0, q(i)), \quad \|q\|_\infty^- = \max_{i \in N} \max(0, -q(i)).$$

Note that

$$\|q\|_\infty = \max(\|q\|_\infty^+, \|q\|_\infty^-)$$

holds, and  $\|q\|_\infty^+ + \|q\|_\infty^-$  serves as a norm of  $q$  (satisfying the axioms of norms). Accordingly, the value  $\|p^* - p\|_\infty^+ + \|p^* - p\|_\infty^-$  represents a distance between two vectors  $p^*$  and  $p$ . For  $p \in \mathbb{Z}^n$ , we define

$$\mu(p) = \min\{\|p^* - p\|_\infty^+ + \|p^* - p\|_\infty^- \mid p^* \in \arg \min g\},$$

which measures the distance between the vector  $p$  and a minimizer of  $g$ .

It is easy to see that  $\mu(p)$  remains the same or decreases by one if  $p$  is updated by adding or subtracting a 0-1 vector. This implies that  $\mu(p) + 1$  is a lower bound for the number of iterations in STEEPESTDESCENT. This is also an upper bound as follows.

**Theorem 1.2.** *The algorithm STEEPESTDESCENT terminates exactly in  $\mu(p^\circ) + 1$  iterations.*

We also consider the following variant of the steepest descent algorithm, where the vector  $p$  is always incremented.

**Algorithm STEEPESTDESCENTUP**

Step0: Set  $p := p^\circ$ , where  $p^\circ \in \mathbb{Z}^n$  is a lower bound of some  $p^* \in \arg \min g$ .

Step1: Find  $X \subseteq N$  that minimizes  $g(p + \chi_X)$ .

Step2: If  $g(p + \chi_X) = g(p)$ , then output  $p$  and stop.

Step3: Set  $p := p + \chi_X$  and go to Step 1.

For the analysis of STEEPESTDESCENTUP, we define

$$\hat{\mu}(p) = \min\{\|p^* - p\|_\infty \mid p^* \in \arg \min g, p^* \geq p\} \quad (p \in \mathbb{Z}^n).$$

**Theorem 1.3.** *Suppose that the initial vector  $p^\circ \in \text{dom } g$  in the algorithm STEEPESTDESCENTUP is a lower bound of some minimizer of  $g$ . Then, the algorithm outputs a minimizer of  $g$  and terminates exactly in  $\hat{\mu}(p^\circ) + 1$  iterations.*

Similarly to STEEPESTDESCENTUP, we can consider an algorithm STEEPESTDESCENTDOWN, where the vector is decreased by a vector  $\chi_X \in \{0, 1\}^n$  that minimizes  $g(p - \chi_X)$ . We define

$$\check{\mu}(p) = \min\{\|p^* - p\|_\infty \mid p^* \in \arg \min g, p^* \leq p\} \quad (p \in \mathbb{Z}^n).$$

**Theorem 1.4.** *Suppose that the initial vector  $p^\circ \in \text{dom } g$  in the algorithm STEEPESTDESCENTDOWN is an upper bound of some minimizer of  $g$ . Then, the algorithm outputs a minimizer of  $g$  and terminates exactly in  $\check{\mu}(p^\circ) + 1$  iterations.*

Theorems 1.2, 1.3, and 1.4 show that the trajectory of a vector  $p$  generated by the steepest descent algorithms is the “shortest” path between the initial vector and a minimizer of  $g$ . This reveals an additional advantage of the steepest descent algorithms, which is important in applications such as iterative auction and computer vision. The proofs of Theorems 1.2 and 1.3 are given in Section 3. The proof of Theorem 1.4 is essentially the same as that for Theorem 1.3 and omitted.




Figure 1: Behavior of steepest descent algorithms. The number associated with each integral lattice point shows the function value of  $g$  at that point. The shaded region shows the set of minimizers of  $g$ .

**An Example** We illustrate the behavior of the algorithms STEEPESTDSCENT and STEEPESTDSCENTUP for an  $L^1$ -convex function  $g : \mathbb{Z}^2 \rightarrow \mathbb{R} \cup \{+\infty\}$  given by

$$\begin{aligned} \text{dom } g &= \{(p(1), p(2)) \in \mathbb{Z}^2 \mid 0 \leq p(i) \leq 4 \ (i = 1, 2)\}, \\ g(p(1), p(2)) &= \max(0, -p(1) + 2, -p(2) + 1, \\ &\quad -p(1) + p(2) - 1, p(1) - p(2) - 2) \\ &\quad \quad \quad ((p(1), p(2)) \in \text{dom } g) \end{aligned}$$

(see Figure 1).

If we apply STEEPESTDSCENT with the initial vector  $p^\circ = (1, 4)$ , then the trajectory of vector  $p$  is one of the three paths depicted by dotted arrows in Figure 1 and the minimizer found by the algorithm is either  $(3, 4)$  or  $(2, 3)$ . Note that  $\mu(p^\circ) = 2$  with

$$\|p^* - p^\circ\|_\infty^+ + \|p^* - p^\circ\|_\infty^- = \begin{cases} 2 + 0 & \text{for } p^* = (3, 4), \\ 1 + 1 & \text{for } p^* = (2, 3). \end{cases}$$

On the other hand, if we apply STEEPESTDSCENT (or STEEPESTDSCENTUP) with the initial vector  $p^\circ = (0, 0)$ , then the vector  $p$  follows one of the three paths depicted by dotted arrows and the minimizer found by the algorithm is either  $(2, 1)$  or  $(2, 2)$ . We have  $\mu(p^\circ) = 2$  with

$$\|p^* - p^\circ\|_\infty^+ + \|p^* - p^\circ\|_\infty^- = 2 + 0 \quad \text{for } p^* = (2, 1), (2, 2),$$

and

$$\hat{\mu}(p^\circ) = \|p^* - p^\circ\|_\infty = 2 \quad \text{for } p^* = (2, 1), (2, 2).$$

## 2 Examples of $L^{\natural}$ -convex Functions and Minimization Algorithms

In this section we show some examples of  $L^{\natural}$ -convex functions arising from applications. We also point out that the minimization algorithms used in those applications can be regarded as special cases of the steepest descent algorithms for  $L^{\natural}$ -convex functions.

### 2.1 Hassin's Algorithm for Minimum Cost Flow Problem

For a directed graph  $G = (V, E)$ , nonnegative edge capacity  $c(e)$ , and edge cost  $\gamma(e) \in \mathbb{R}$  for  $e \in E$ , the minimum cost flow problem is formulated as:

$$\begin{aligned} \text{Minimize} \quad & \sum_{(u,v) \in E} \gamma(u,v)x(u,v) \\ \text{subject to} \quad & \sum_{v:(u,v) \in E} x(u,v) - \sum_{v:(v,u) \in E} x(v,u) = 0 \quad (u \in V), \\ & 0 \leq x(u,v) \leq c(u,v) \quad ((u,v) \in E). \end{aligned}$$

The dual problem is given as:

$$\begin{aligned} \text{Maximize} \quad & g_{\text{H}}(p) \equiv \sum_{(u,v) \in E} c(u,v) \min\{0, p(u) - p(v) + \gamma(u,v)\} \\ \text{subject to} \quad & p(v) \in \mathbb{R} \quad (v \in V). \end{aligned}$$

We here assume that edge cost  $\gamma(u,v)$  is integer-valued. Then, there exists an integral optimal solution to the dual problem, and we may assume that  $p(v) \in \mathbb{Z}$  ( $v \in V$ ) in the dual problem.

It is known that  $g_{\text{H}}$  is an  $L^{\natural}$ -concave function ( $-g_{\text{H}}$  is  $L^{\natural}$ -convex) if we regard  $g_{\text{H}}$  as a function in integer vectors (see [14]). Since  $g_{\text{H}}(p + \mathbf{1}) = g_{\text{H}}(p)$  ( $\forall p \in \mathbb{Z}^V$ ), for each  $p^{\circ} \in \mathbb{Z}^V$ , there exists an optimal solution  $p^* \in \mathbb{Z}^V$  satisfying  $p^* \geq p^{\circ}$ .

Hassin's algorithm in [10] can be seen as an application of algorithm STEEPESTDESCENTUP to the  $L^{\natural}$ -convex function  $-g_{\text{H}}$ ; see [16] for details. We also mention that the algorithm by Chung and Tcha [5] for the minimum-cost submodular flow problem, which is a generalization of Hassin's algorithm, can also be seen as a special implementation of algorithm STEEPESTDESCENTUP.

### 2.2 Discrete Optimization Approach in Computer Vision

Given an undirected graph  $G = (V, E)$  and univariate convex functions  $\varphi_u : \mathbb{Z} \rightarrow \mathbb{R} \cup \{+\infty\}$  ( $u \in V$ ) and  $\psi_{uv} : \mathbb{Z} \rightarrow \mathbb{R} \cup \{+\infty\}$  ( $(u,v) \in E$ ),

consider the following optimization problem:

$$\begin{aligned} \text{(P): Minimize } & g_{\text{CV}}(p) \equiv \sum_{u \in V} \varphi_u(p(u)) + \sum_{(u,v) \in E} \psi_{uv}(p(v) - p(u)) \\ \text{subject to } & p \in \mathbb{Z}^V. \end{aligned}$$

It is known that the objective function  $g_{\text{CV}}$  of this problem is an  $L^{\natural}$ -convex function (see [13, 14]).

The problem (P) arises in many applications in computer vision such as panoramic image stitching [20], image restoration [4], minimization of total variation [7], and phase unwrapping in SAR images [2]. In such applications, the node set  $V$  of the undirected graph  $G = (V, E)$  usually corresponds to the set of pixels in a given image, and variable  $p(u)$  is the “label” of pixel  $u \in V$  that represents disparity, intensity, etc. Functions  $\varphi_u$  encode unary data penalty functions, and  $\psi_{uv}$  are pairwise interaction potentials. The objective function of (P) is often derived in the context of Markov random fields [8]; a minimizer of the function  $g_{\text{CV}}$  corresponds to a maximum a-posteriori labeling.

There have been proposed many algorithms for (P) in computer vision (see, e.g., [2, 13]). Among them, the primal algorithm of Kolmogorov and Shioura [13] can be seen as an application of algorithm `STEEPESTDESCENT` to (P), while the algorithm of Bioucas-Dias and Valadão [2] is an application of `STEEPESTDESCENTUP` to a special case of (P) where the term  $\sum_{u \in V} \varphi_u(p(u))$  in the objective function is missing.

### 2.3 Iterative Auction in Mathematical Economics

In an auction, we want to find “good” prices for items to be allocated to bidders. An algorithm for computing such “good” prices, called ascending auction [1], can be seen as a special implementation of algorithm `STEEPESTDESCENTUP`.

We consider an auction market with  $n$  types of items or goods, denoted by  $N = \{1, 2, \dots, n\}$ , and  $m$  bidders, denoted by  $M = \{1, 2, \dots, m\}$ . Each bidder  $j \in M$  has his valuation function  $f_j : 2^N \rightarrow \mathbb{R}$ ; the value  $f_j(X)$  represents the degree of satisfaction for an item set  $X \subseteq N$ . We assume that each  $f_j$  satisfies the so-called “gross-substitutes” condition, which is a natural assumption for valuation functions (see [1, 9, 12] for a precise definition). We also assume that each  $f_j$  is integer-valued function. An allocation of items is defined as a family of item sets  $X_1, X_2, \dots, X_m$  satisfying  $X_j \cap X_k = \emptyset$  if  $j \neq k$  and  $\bigcup_{j \in M} X_j = N$ .

Given a price vector  $p \in \mathbb{R}^n$ , each bidder  $j \in M$  wants to have an item set  $X$  which maximizes the value  $f_j(X) - p(X)$ , where  $p(X) = \sum_{i \in X} p(i)$ . On the other hand, the auctioneer wants to find a price vector under which all items are sold completely. Hence, all of the auctioneer and bidders are happy



if we can find a pair of a price vector  $p^*$  and an allocation  $X_1^*, X_2^*, \dots, X_m^*$  satisfying the condition

$$X_j^* \in \arg \max \{f_j(X) - p(X) \mid X \subseteq N\} \quad (j \in M).$$

Such a pair is called a Walrasian equilibrium;  $p^*$  is called a Walrasian equilibrium price vector (see, e.g., [3, 6]).

In the auction literature an algorithm called the iterative auction (or dynamic auction, Walrasian tâtonnement process, etc.) is often used to find an equilibrium [3, 6]. An iterative auction finds an equilibrium price vector by iteratively updating a current price vector  $p$ . The most natural and popular iterative auction is the ascending auction, in which the current price vector is increased monotonically. The ascending auction is a natural generalization of the classical English auction for a single item, and known to have various nice properties (see, e.g., [3, 6]); in particular, it is natural from the economic point of view, and easy to understand and implement.

The ascending auction presented in Ausubel [1] uses a function defined by

$$L(p) = \sum_{j=1}^m \max \{f_j(X) - p(X) \mid X \subseteq N\} + p(N) \quad (p \in \mathbb{R}^n),$$

which is called the Lyapunov function. Under the assumption that each  $f_j$  satisfies the gross substitutes condition,  $p^*$  is an equilibrium price vector if and only if it is a minimizer of the Lyapunov function and that there exists an integral minimizer  $p^* \in \mathbb{Z}^n$  of the Lyapunov function. Based on this fact, the ascending auction in [1] tries to find a minimizer of the Lyapunov function.

It can be shown that the Lyapunov function  $L$  is an  $L^\natural$ -concave function if it is regarded as a function in integer vectors, which follows from the conjugacy results in discrete convex analysis and the assumption that each  $f_j$  satisfies the gross substitutes condition (see, e.g., [14, 18]). Moreover, it is observed in [18] that the ascending auction in [1] can be seen as an application of algorithm STEEPESTDESCENTUP to the function  $-L$ .

### 3 Proofs

In this section, we prove Theorems 1.2 and 1.3. The key fact used in our proofs is the following property of  $L^\natural$ -convex functions. For  $p \in \mathbb{Z}^n$ , we denote  $\text{supp}^+(p) = \{i \in N \mid p(i) > 0\}$ .

**Lemma 3.1** ([14, Th. 7.7]). *Let  $g : \mathbb{Z}^n \rightarrow \mathbb{R} \cup \{+\infty\}$  be an  $L^\natural$ -convex function. For every  $p, q \in \text{dom } g$  with  $\text{supp}^+(p - q) \neq \emptyset$ , it holds that*

$$g(p) + g(q) \geq g(p - \chi_X) + g(q + \chi_X),$$

where  $X = \arg \max_{i \in N} \{p(i) - q(i)\}$ .

### 3.1 Proof of Theorem 1.2

The bound  $\mu(p^\circ) + 1$  for the number of iterations in algorithm STEEPEST-DESCENT can be obtained by repeated application of the following lemma.

**Lemma 3.2.** *Let  $p \in \mathbb{Z}^n$  be a vector with  $\mu(p) > 0$ . Suppose that  $\sigma \in \{+1, -1\}$  and  $X \subseteq N$  minimize the value  $g(p + \sigma\chi_X)$ . Then,  $\mu(p + \sigma\chi_X) = \mu(p) - 1$ .*

*Proof.* We consider the case with  $\sigma = +1$  since the other case with  $\sigma = -1$  can be dealt with similarly. For every  $d \in \mathbb{Z}^n$  and  $Y \subseteq N$ , we have

$$\|d - \chi_Y\|_\infty^+ \geq \|d\|_\infty^+ - 1, \quad \|d - \chi_Y\|_\infty^- \geq \|d\|_\infty^-.$$

Hence, it holds that

$$\begin{aligned} & \mu(p + \chi_X) \\ &= \min\{\|q - (p + \chi_X)\|_\infty^+ + \|q - (p + \chi_X)\|_\infty^- \mid q \in \arg \min g\} \\ &\geq \min\{\|q - p\|_\infty^+ + \|q - p\|_\infty^- \mid q \in \arg \min g\} - 1 \\ &= \mu(p) - 1. \end{aligned}$$

In the following, we prove that the reverse inequality  $\mu(p + \chi_X) \leq \mu(p) - 1$  holds.

We denote

$$S = \{q \in \arg \min g \mid \|q - p\|_\infty^+ + \|q - p\|_\infty^- = \mu(p)\}.$$

Let  $q^*$  be a vector in  $S$  with the maximum value of  $\|q^* - p\|_\infty^+$ .

**Claim 1:**  $\|q^* - p\|_\infty^+ > 0$ .

[Proof of Claim 1] Assume, to the contrary, that  $q^* \leq p$ . Note that this assumption implies  $\|q^* - p\|_\infty^- > 0$  since  $\mu(p) > 0$ .

By  $L^1$ -convexity of  $g$  in (1), we have

$$\begin{aligned} & g(p + \chi_X) + g(q^*) \\ &\geq g((p + \chi_X - \mathbf{1}) \vee q^*) + g((p + \chi_X) \wedge (q^* + \mathbf{1})). \end{aligned} \quad (2)$$

Let  $Z = \{i \in N \mid q^*(i) - p(i) = 0\}$ , which may be the empty set. Then, we have

$$(p + \chi_X - \mathbf{1}) \vee q^* = p - \chi_{N \setminus (X \cup Z)}, \quad (p + \chi_X) \wedge (q^* + \mathbf{1}) = q^* + \chi_{(N \setminus Z) \cup X},$$

which, together with (2), implies

$$g(p + \chi_X) + g(q^*) \geq g(p - \chi_{N \setminus (X \cup Z)}) + g(q^* + \chi_{(N \setminus Z) \cup X}). \quad (3)$$

By the choice of  $\sigma = +1$  and  $X$ , we have  $g(p + \chi_X) \leq g(p - \chi_{N \setminus (X \cup Z)})$ . From this and (3) follows that  $g(q^*) \geq g(q^* + \chi_{(N \setminus Z) \cup X})$ , implying that  $q^* + \chi_{(N \setminus Z) \cup X} \in \arg \min g$ . We also have

$$\begin{aligned} \|(q^* + \chi_{(N \setminus Z) \cup X}) - p\|_\infty^- &= \max_{i \in N \setminus Z} \{p(i) - (q^*(i) + 1)\} \\ &= \|q^* - p\|_\infty^- - 1, \\ \|(q^* + \chi_{(N \setminus Z) \cup X}) - p\|_\infty^+ &\leq 1 = \|q^* - p\|_\infty^+ + 1 \end{aligned} \quad (4)$$

since  $\|q^* - p\|_\infty^- > 0$  and  $\|q^* - p\|_\infty^+ = 0$ . Hence, it follows that

$$\begin{aligned} \|(q^* + \chi_{(N \setminus Z) \cup X}) - p\|_\infty^+ + \|(q^* + \chi_{(N \setminus Z) \cup X}) - p\|_\infty^- \\ \leq \|q^* - p\|_\infty^+ + \|q^* - p\|_\infty^- = \mu(p). \end{aligned}$$

By the definition of  $\mu(p)$ , this inequality and the inequality (4) must hold with equality. Hence, we have  $q^* + \chi_{(N \setminus Z) \cup X} \in S$  and

$$\|(q^* + \chi_{(N \setminus Z) \cup X}) - p\|_\infty^+ = \|q^* - p\|_\infty^+ + 1 > \|q^* - p\|_\infty^+.$$

This, however, is a contradiction to the choice of  $q^*$ .

[End of Proof of Claim 1]

In the following, we further assume that  $q^*$  is a minimal vector in the set  $S$  with  $\|q^* - p\|_\infty^+ = \xi$ , where

$$\xi = \max\{\|q - p\|_\infty^+ \mid q \in S\}.$$

Note that  $\xi > 0$  by Claim 1.

We denote

$$A = \arg \max_{i \in N} \{q^*(i) - p(i)\}, \quad B = \arg \min_{i \in N} \{q^*(i) - p(i)\}.$$

**Claim 2:** We have  $A \subseteq X$ .

[Proof of Claim 2] Assume, to the contrary, that  $A \setminus X \neq \emptyset$  holds. We claim that  $q^* - \chi_{A \setminus X} \in \arg \min g$ . Since  $\|q^* - p\|_\infty^+ = \xi > 0$ , we have  $A \subseteq \text{supp}^+(q^* - p)$ , from which follows that

$$\text{supp}^+(q^* - (p + \chi_X)) \supseteq A \setminus X \neq \emptyset.$$

We also have

$$\arg \max_{i \in N} \{q^*(i) - (p + \chi_X)(i)\} = A \setminus X.$$

Hence, Lemma 3.1 implies that

$$\begin{aligned} g(q^*) + g(p + \chi_X) &\geq g(q^* - \chi_{A \setminus X}) + g(p + \chi_X + \chi_{A \setminus X}) \\ &= g(q^* - \chi_{A \setminus X}) + g(p + \chi_{X \cup A}). \end{aligned} \quad (5)$$

By the choice of  $X$ , we have  $g(p + \chi_X) \leq g(p + \chi_{X \cup A})$ , which, together with (5), implies that  $g(q^*) \geq g(q^* - \chi_{A \setminus X})$ , i.e.,  $q^* - \chi_{A \setminus X} \in \arg \min g$ .

Since  $A \setminus X \subseteq \text{supp}^+(q^* - p)$ , we have

$$\begin{aligned} \|(q^* - \chi_{A \setminus X}) - p\|_\infty^+ &\leq \|q^* - p\|_\infty^+, \\ \|(q^* - \chi_{A \setminus X}) - p\|_\infty^- &= \|q^* - p\|_\infty^-, \end{aligned} \quad (6)$$

from which follows that

$$\|(q^* - \chi_{A \setminus X}) - p\|_\infty^+ + \|(q^* - \chi_{A \setminus X}) - p\|_\infty^- \leq \|q^* - p\|_\infty^+ + \|q^* - p\|_\infty^- = \mu(p).$$

By the definition of  $\mu(p)$ , this inequality and the inequality (6) must hold with equality. Hence, the vector  $q^* - \chi_{A \setminus X}$  belongs to  $S$  with  $\|(q^* - \chi_{A \setminus X}) - p\|_\infty^+ = \xi$ , a contradiction to the minimality of  $q^*$ . [End of Proof of Claim 2]

To show the inequality  $\mu(p + \chi_X) \leq \mu(p) - 1$ , we first consider the case with  $\min_{i \in N} \{q^*(i) - p(i)\} > 0$ . Then, we have

$$\|q^* - (p + \chi_X)\|_\infty^- = 0 = \|q^* - p\|_\infty^-$$

since  $q^* \geq p + \chi_X$ . Claim 2 implies that

$$\|q^* - (p + \chi_X)\|_\infty^+ = \|q^* - p\|_\infty^+ - 1.$$

Therefore, it follows that

$$\begin{aligned} \mu(p + \chi_X) &\leq \|q^* - (p + \chi_X)\|_\infty^+ + \|q^* - (p + \chi_X)\|_\infty^- \\ &= (\|q^* - p\|_\infty^+ - 1) + \|q^* - p\|_\infty^- = \mu(p) - 1. \end{aligned}$$

We next consider the case with  $\min_{i \in N} \{q^*(i) - p(i)\} \leq 0$ . We claim that  $q^* + \chi_{B \cap X} \in \arg \min g$  holds. If  $B \cap X = \emptyset$ , then  $q^* + \chi_{B \cap X} = q^* \in \arg \min g$ . Hence, we assume  $B \cap X \neq \emptyset$ . Since

$$\text{supp}^+((p + \chi_X) - q^*) \neq \emptyset, \quad \arg \max_{i \in N} \{(p + \chi_X)(i) - q^*(i)\} = B \cap X,$$

it follows from Lemma 3.1 that

$$\begin{aligned} g(p + \chi_X) + g(q^*) &\geq g(p + \chi_X - \chi_{B \cap X}) + g(q^* + \chi_{B \cap X}) \\ &= g(p + \chi_{X \setminus B}) + g(q^* + \chi_{B \cap X}). \end{aligned} \quad (7)$$

By the choice of  $X$ , we have  $g(p + \chi_X) \leq g(p + \chi_{X \setminus B})$ , which, together with (7), implies that  $g(q^*) \geq g(q^* + \chi_{B \cap X})$ , i.e.,  $q^* + \chi_{B \cap X}$  is also a minimizer of  $g$ .

Since  $\min_{i \in N} \{q^*(i) - p(i)\} \leq 0 < \max_{i \in N} \{q^*(i) - p(i)\}$ , we have  $A \cap B = \emptyset$ , which, together with Claim 2, implies  $A \subseteq X \setminus B$ . Hence, it holds that

$$\|(q^* + \chi_{B \cap X}) - (p + \chi_X)\|_\infty^+ = \|q^* - p - \chi_{X \setminus B}\|_\infty^+ = \|q^* - p\|_\infty^+ - 1.$$

We also have

$$\|(q^* + \chi_{B \cap X}) - (p + \chi_X)\|_\infty^- = \|q^* - p - \chi_{X \setminus B}\|_\infty^- = \|q^* - p\|_\infty^-,$$

where the second equality follows from the definition of  $B$ . Hence, it holds that

$$\begin{aligned} & \mu(p + \chi_X) \\ & \leq \|(q^* + \chi_{B \cap X}) - (p + \chi_X)\|_\infty^+ + \|(q^* + \chi_{B \cap X}) - (p + \chi_X)\|_\infty^- \\ & = (\|q^* - p\|_\infty^+ - 1) + \|q^* - p\|_\infty^- = \mu(p) - 1. \end{aligned}$$

□

### 3.2 Proof of Theorem 1.3

The proof of Theorem 1.3 is quite similar to and simpler than that of Theorem 1.2. Theorem 1.3 can be proved by using the following property repeatedly.

**Lemma 3.3.** *Let  $p \in \mathbb{Z}^n$  be a vector with  $\hat{\mu}(p) > 0$ , and  $X \subseteq N$  be a set that minimizes the value of  $g(p + \chi_X)$ . Then,  $\hat{\mu}(p + \chi_X) = \hat{\mu}(p) - 1$ .*

*Proof.* The inequality  $\hat{\mu}(p + \chi_X) \geq \hat{\mu}(p) - 1$  can be shown as follows. By the triangle inequality, we have  $\|q - (p + \chi_X)\|_\infty \geq \|q - p\|_\infty - 1$  for every  $q \in \mathbb{Z}^n$ . Taking the minimum over all  $q \in \arg \min g$  with  $q \geq p + \chi_X$ , we obtain

$$\begin{aligned} & \hat{\mu}(p + \chi_X) \\ & \geq \min\{\|q - p\|_\infty \mid q \in \arg \min g, q \geq p + \chi_X\} - 1 \\ & \geq \min\{\|q - p\|_\infty \mid q \in \arg \min g, q \geq p\} - 1 = \hat{\mu}(p) - 1. \end{aligned}$$

In the following, we show that the reverse inequality  $\hat{\mu}(p + \chi_X) \leq \hat{\mu}(p) - 1$  holds.

Let  $p^*$  be a vector such that  $p^* \in \arg \min g$ ,  $p^* \geq p$ , and  $\|p^* - p\|_\infty = \hat{\mu}(p)$ , and assume that  $p^*$  is minimal among such vectors. We denote

$$A = \arg \max_{i \in N} \{p^*(i) - p(i)\}.$$

We have  $p^* \neq p$  since  $\|p^* - p\|_\infty = \hat{\mu}(p) > 0$ .

**Claim 1:** We have  $A \subseteq X$ .

[Proof of Claim 1] Assume, to the contrary, that  $A \setminus X \neq \emptyset$  holds. Since  $A \subseteq \text{supp}^+(p^* - p)$ , we have

$$\text{supp}^+(p^* - (p + \chi_X)) \supseteq A \setminus X \neq \emptyset.$$

We also have

$$\arg \max_{i \in N} \{p^*(i) - (p + \chi_X)(i)\} = A \setminus X.$$

Hence, Lemma 3.1 implies that

$$\begin{aligned} g(p^*) + g(p + \chi_X) &\geq g(p^* - \chi_{A \setminus X}) + g(p + \chi_X + \chi_{A \setminus X}) \\ &= g(p^* - \chi_{A \setminus X}) + g(p + \chi_{X \cup A}). \end{aligned} \quad (8)$$

Since  $p^* \geq p^* - \chi_{A \setminus X} \geq p$ , we have  $g(p^*) < g(p^* - \chi_{A \setminus X})$  by the choice of  $p^*$ . This inequality, together with (8), implies that  $g(p + \chi_X) > g(p + \chi_{X \cup A})$ , a contradiction to the choice of  $X$ . Hence, we have  $A \subseteq X$ . [End of Proof of Claim 1]

Suppose first that the condition  $p^* \geq p + \chi_X$  holds. Then, we have

$$\hat{\mu}(p + \chi_X) \leq \|p^* - (p + \chi_X)\|_\infty = \|p^* - p\|_\infty - 1 = \hat{\mu}(p) - 1,$$

where the first equality is by Claim 1.

We next consider the case where the condition  $p^* \geq p + \chi_X$  fails. Then,  $B \cap X \neq \emptyset$  for  $B = \{i \in N \mid p^*(i) = p(i)\}$ . Since  $p^* \geq p$ , we have

$$p^*(i) = p(i) \quad (\forall i \in B), \quad p^*(i) > p(i) \quad (\forall i \in N \setminus B), \quad (9)$$

from which  $p^* + \chi_{B \cap X} \geq p + \chi_X$  follows.

We now show that  $p^* + \chi_{B \cap X} \in \arg \min g$  holds. The condition (9) implies

$$\text{supp}^+((p + \chi_X) - p^*) = \arg \max_{i \in N} \{(p + \chi_X)(i) - p^*(i)\} = B \cap X.$$

Hence, it follows from Lemma 3.1 that

$$\begin{aligned} g(p + \chi_X) + g(p^*) &\geq g(p + \chi_X - \chi_{B \cap X}) + g(p^* + \chi_{B \cap X}) \\ &= g(p + \chi_{X \setminus B}) + g(p^* + \chi_{B \cap X}). \end{aligned} \quad (10)$$

By the choice of  $X$ , we have  $g(p + \chi_X) \leq g(p + \chi_{X \setminus B})$ , which, together with (10), implies that  $g(p^*) \geq g(p^* + \chi_{B \cap X})$ , i.e.,  $p^* + \chi_{B \cap X}$  is also a minimizer of  $g$ .

Since  $A \subseteq X \setminus B$  by Claim 1 and the definitions of  $A$  and  $B$ , it holds that

$$\begin{aligned} \hat{\mu}(p + \chi_X) &\leq \|(p^* + \chi_{B \cap X}) - (p + \chi_X)\|_\infty \\ &= \|p^* - p - \chi_{X \setminus B}\|_\infty \\ &= \|p^* - p\|_\infty - 1 = \hat{\mu}(p) - 1. \end{aligned}$$

□

## 4 Conclusion

The concept of  $L^1$ -convexity is generalized to polyhedral convex functions. A steepest descent algorithm similar to STEEPESTDESCENT works for the minimization of a polyhedral  $L^1$ -convex function, and has a similar property as STEEPESTDESCENT; in particular, the trajectory of a vector  $p$  generated by the algorithm is the “shortest” path between the initial vector and a minimizer (see [17] for details).

Part of the results in this paper is presented (without proofs) in the extended abstract [18] in Proceedings of 24th International Symposium on Algorithms and Computation (ISAAC 2013). This research is supported by KAKENHI (21360045, 21740060, 24500002) and the Aihara Project, the FIRST program from JSPS.

## References

- [1] L. M. Ausubel, An efficient dynamic auction for heterogeneous commodities, *Amer. Econ. Rev.* 96 (2006) 602–629.
- [2] J. M. Bioucas-Dias, G. Valadão, Phase unwrapping via graph cuts, *IEEE Trans. Image Process.* 16 (2007) 698–709.
- [3] L. Blumrosen, N. Nisan, Combinatorial auction. in: N. Nisan, et al. (eds.) *Algorithmic Game Theory*, Cambridge Univ. Press, Cambridge, 2007, pp. 267–299.
- [4] Y. B. Boykov, O. Veksler, R. Zabih, Fast approximate energy minimization via graph cuts, *IEEE Trans. Pattern Anal. Mach. Intell.* 23 (2001) 1222–1239.
- [5] N.-k. Chung, D.-w. Tcha, A dual algorithm for submodular flow problems, *Oper. Res. Lett.* 10 (1991) 489–495.
- [6] P. Cramton, Y. Shoham, R. Steinberg, *Combinatorial Auctions*. MIT Press, Cambridge, MA, 2006.
- [7] J. Darbon, M. Sigelle, Image restoration with discrete constrained total variation, part I: fast and exact optimization, *J. Math. Imaging Vision* 26 (2006) 261–276.
- [8] S. Geman, D. Geman, Stochastic relaxation, Gibbs distributions, and the Bayesian restoration of images, *IEEE Trans. Pattern Anal. Mach. Intell.* 6 (1984) 721–741.
- [9] F. Gul, E. Stacchetti, The English auction with differentiated commodities, *J. Economic Theory* 92 (2000) 66–95.

- [10] R. Hassin, The minimum cost flow problem: a unifying approach to dual algorithms and a new tree-search algorithm, *Math. Program.* 25 (1983) 228–239.
- [11] W.T. Huh, G. Janakiraman, On the optimal policy structure in serial inventory systems with lost sales, *Oper. Res.* 58 (2010) 486–491.
- [12] A. S. Kelso, V. P. Crawford, Job matching, coalition formation and gross substitutes, *Econometrica* 50 (1982) 1483–1504.
- [13] V. Kolmogorov, A. Shioura, New algorithms for convex cost tension problem with application to computer vision, *Discrete Optim.* 6 (2009) 378–393.
- [14] K. Murota, *Discrete Convex Analysis*, SIAM, Philadelphia, PA, 2003.
- [15] K. Murota, On steepest descent algorithms for discrete convex functions, *SIAM J. Optim.* 14 (2003) 699–707.
- [16] K. Murota, A. Shioura, Dijkstra’s algorithm and L-concave function maximization, *Math. Program.* (2013) published online.
- [17] K. Murota, A. Shioura, On some properties of polyhedral L-concave maximization algorithm, Technical report METR 2013-18, Department of Mathematical Informatics, University of Tokyo (2013).
- [18] K. Murota, A. Shioura, Z. Yang, Computing a Walrasian equilibrium in iterative auctions with multiple differentiated items, in: *Proc. 24th International Symposium on Algorithms and Computation (ISAAC 2013)*, to appear.
- [19] P. Zipkin, On the structure of lost-sales inventory models, *Oper. Res.* 56 (2008) 937–944.
- [20] A. Zomet, A. Levin, S. Peleg, Y. Weiss, Seamless image stitching by minimizing false edges, *IEEE Trans. Image Process.* 15 (2006) 969–977.